

MATH 545, Stochastic Calculus

Problem set 1

January 15, 2020

These problems are due on MON Jan 27. You can give them to me in class, drop them in my box. In all of the problems \mathbb{E} denotes the expected value with respect to the specified probability measure \mathbb{P} .

Problem 0. Read [Klebaner] Chapter 3 (by JAN 29)

Problem 1. Consider the following probability space $(\Omega, \mathcal{F}, \mathbb{P})$ where $\Omega = \{(\omega_0, \omega_1) : \omega_i \in \{-1, 0, 1\}\}$. Take of each of the ω_i to be mutually independent with $\mathbb{P}[\omega_i = 0] = 1/2$ and $\mathbb{P}[\omega_i = \pm 1] = 1/4$. (\mathcal{F} is just the σ -algebra generated by the collection of single points, but this is not important).

For $n = 0$ or 1 define the random variables X_n by $X_0 = \omega_0$ and $X_1 = \omega_0\omega_1$.

- What is $\mathbb{P}[X_1 = 1]$? What is $\mathbb{E}[X_0]$?
- Let A be the event that $\{X_1 \neq 0\}$. What is $\sigma(A)$?
- What is $\sigma(X_0)$ and $\sigma(X_1)$?
- What is $\mathbb{E}[X_1|A]$? What is $\mathbb{E}[X_1|X_0]$?

Problem 2. Consider the sample space $\Omega = \{\omega_i\}_{i=1}^3$ with ω_i are independent identically distributed according to

$$\omega_i = \begin{cases} 1 & \text{with probability } 1/2 \\ -1 & \text{with probability } 1/2 \end{cases}$$

Furthermore, define $X_n(\omega) = \sum_{i=1}^n \omega_i$ and let $\mathcal{F}_1 = \sigma(X_1)$, $\mathcal{F}_{12} = \sigma(\{\omega_1, \omega_2\})$ (the sigma algebras generated by the outcome of the first flip and of the first two flips respectively). Compute

- $\mathbb{E}[X_3|X_1]$
- $\mathbb{E}[\mathbb{E}[X_3|\mathcal{F}_1]|\mathcal{F}_{12}]$
- $\mathbb{E}[\mathbb{E}[X_3|\mathcal{F}_{12}]|\mathcal{F}_1]$
- check that the answer to the above two points is the same (tower property)

Suggestion: write a random variable Y (conditional expectations are random variables!) on σ -algebra \mathcal{F} with a countable number of elements as

$$Y(\omega) = \sum_k 1_{A_k}(\omega) a_k \quad \text{for some } A_k \in \mathcal{F}, a_k \in \mathbb{R},$$

where $1_A(\omega)$ is the indicator function on the set A .

Problem 3. For a general probability space $(\Omega, \mathcal{F}, \mathbb{P})$, prove the Chebychev inequality:

$$\mathbb{P}[|X| \geq \lambda] \leq \frac{\mathbb{E}[|X|^p]}{\lambda^p}$$

for all $1 \leq p < \infty$ as well as its exponential form

$$\mathbb{P}[|X| \geq \lambda] \leq \frac{\mathbb{E}[e^{k|X|}]}{e^{k\lambda}}$$

when the integrals on the right are finite. Here k and λ are positive constants. (Hint: denoting by $\mathbb{1}\{A\}$ the indicator function on the set A combine the facts that $\mathbb{P}[|X| \geq \lambda] = \mathbb{E}[\mathbb{1}\{|X| \geq \lambda\}]$, $\mathbb{1}\{|X| \geq \lambda\} \lambda^p \leq \mathbb{1}\{|X| \geq \lambda\} |X|^p$, and $1 = \mathbb{1}\{|X| \geq \lambda\} + \mathbb{1}\{|X| < \lambda\}$.)

Problem 4. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $A_k, k = 1, 2, 3, \dots$ be a sequence of events in \mathcal{F} for which

$$\sum_{k=1}^{\infty} \mathbb{P}[A_k] < \infty$$

Prove the (first) Borell-Cantelli lemma which says that

$$\mathbb{P}\left[\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} A_k\right] = 0$$

Interpret this event whose probability is zero. (Say something like “it is the event in which all of the events A_k happen”. This of course not the right answer.)

Now let us use this result: Let X_k be a sequence of random variables and suppose that there is another random variable X such that if

$$A_k = \{\omega \in \Omega \mid |X_k(\omega) - X(\omega)| > \delta\}$$

then

$$\sum_{k=1}^{\infty} \mathbb{P}[A_k] < \infty$$

for all $\delta > 0$. Show that $X_n \rightarrow X$ with probability one. (That is the set of ω so that $X_n(\omega) \rightarrow X(\omega)$ has measure (probability) one.)

Basically, the first Borell-Cantelli lemma is the principle means by which information about expectations can be converted into almost sure information.

Problem 5 (optional). If $\mathcal{F}_i, i \in I$ for some index set I , are σ -algebras on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, show that $\mathcal{G} := \bigcap_{i \in I} \mathcal{F}_i$ is a σ -algebra.

Problem 6 (optional). Consider the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ where

$$\Omega = [0, 1], \quad \mathcal{F} = \mathcal{B}_{[0,1]}, \quad \mathbb{P} = \text{Lebesgue measure,}$$

where $\mathcal{B}_{[0,1]}$ denote the Borel σ -algebra on $[0, 1]$. Consider the following two random variables defined on this probability space.

$$X(\omega) = \begin{cases} 1, & \text{if } \omega \in [0, 0.5) \\ 0, & \text{if } \omega \in [0.5, 1] \end{cases} \quad Y(\omega) = \begin{cases} 1, & \text{if } \omega \in [0, 0.25) \cup [0.5, 0.75) \\ 0, & \text{if } \omega \in [0.25, 0.5) \cup [0.75, 1] \end{cases}$$

- Check that X and Y have the same distribution.
- Compute $\mathbb{P}[X = Y]$.
- Write down the sets that are contained in $\sigma(X)$ and $\sigma(Y)$, respectively.
- Show that X and Y are independent.
- Explain how to define a sequence of random variables X_1, X_2, \dots , so that (i) they all have the same distribution as X and (ii) they are mutually independent.

Problem 7 (optional, hard, requires book). Consider a random variable X on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\mathcal{G}_1 \subset \mathcal{F}$ and $\mathcal{G}_2 \subset \mathcal{F}$ be two **independent** sub σ -algebras. Suppose that X is independent of \mathcal{G}_1 . Show that (this is property 5, equation (2.22) on pp. 45 of [Klebaner])

$$\mathbb{E}[X|\mathcal{G}_2] = \mathbb{E}[X|\sigma(\mathcal{G}_1, \mathcal{G}_2)], \quad \text{a.s.}$$

Remark. A definition for the *independence* of two σ -algebras (i.e., σ -fields) is given on page 39 of [Klebaner]. A random variable X is independent of a σ -algebra \mathcal{G} if its generated σ -algebra $\sigma(X)$ is independent of \mathcal{G} .