

EXAM 3

Math 216, 2015-2016 Fall, Clark Bray.

You have 50 minutes.

No notes, no books, no calculators.

YOU MUST SHOW ALL WORK AND EXPLAIN ALL REASONING
TO RECEIVE CREDIT. CLARITY WILL BE CONSIDERED IN GRADING.

All answers must be simplified. All of the policies and guidelines
on the class webpages are in effect on this exam.

Good luck!

Name _____

Disc.: Number _____ TA _____ Day/Time _____

“I have adhered to the Duke Community
Standard in completing this
examination.”

1. _____

2. _____

3. _____

4. _____

5. _____

6. _____

Signature: _____

Total Score _____ (/100 points)

1. (20 pts) The matrix

$$A = \begin{pmatrix} 1 & 2 \\ -3 & 6 \end{pmatrix}$$

has eigenvalues 3 and 4. Find an invertible matrix B and diagonal matrix D with $D = B^{-1}AB$.

2. (20 pts) In the inner product space $C[0, 1]$ using the L^2 inner product, compute the cosine of the angle between the vectors $f(x) = x$ and $g(x) = x^2$.

3. (15 pts) The function $L : V \rightarrow \mathbb{R}^3$ is a linear transformation.

(a) Use L to form a “Wronskian” to help us consider the potential linear independence / dependence of a list of vectors $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ in the vector space V .

(b) Suppose we know the following:

$$L(\vec{v}_1) = \begin{pmatrix} 4 \\ 2 \\ 1 \end{pmatrix} \quad \text{and} \quad L(\vec{v}_2) = \begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix} \quad \text{and} \quad L(\vec{v}_3) = \begin{pmatrix} 0 \\ 1 \\ 9 \end{pmatrix}$$

What can you conclude about the potential linear independence/dependence of the list of vectors $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$?

(c) Suppose we know the following:

$$L(\vec{v}_4) = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} \quad \text{and} \quad L(\vec{v}_5) = \begin{pmatrix} 2 \\ 0 \\ 5 \end{pmatrix} \quad \text{and} \quad L(\vec{v}_6) = \begin{pmatrix} 0 \\ 2 \\ -4 \end{pmatrix}$$

What can you conclude about the potential linear independence/dependence of the list of vectors $\{\vec{v}_4, \vec{v}_5, \vec{v}_6\}$?

4. (15 pts) Suppose we know that $A = B^{-1}CB$, and $\vec{q}'(x) = C\vec{q}(x)$. Find (and justify) a solution to the equation $\vec{y}'(x) = A\vec{y}(x)$.

5. (15 pts) The arithmetic below is given.

$$A = \begin{pmatrix} -11 & -18 \\ 12.5 & 19 \end{pmatrix} = \begin{pmatrix} -3 & 2 \\ 2.5 & -1.5 \end{pmatrix} \begin{pmatrix} 4 & 1 \\ 0 & 4 \end{pmatrix} \begin{pmatrix} 3 & 4 \\ 5 & 6 \end{pmatrix}$$

Find a fundamental set of solutions to the system $\vec{y}' = A\vec{y}$.

6. (15 pts) Find a particular solution to the system below.

$$y_1' = 3y_1 - 2y_2 + 1$$

$$y_2' = 2y_1 + y_2 + e^x$$