

EXAM 3

Math 216, 2015-2016 Spring, Clark Bray.

You have 50 minutes.

No notes, no books, no calculators.

YOU MUST SHOW ALL WORK AND EXPLAIN ALL REASONING
TO RECEIVE CREDIT. CLARITY WILL BE CONSIDERED IN GRADING.

All answers must be simplified. All of the policies and guidelines
on the class webpages are in effect on this exam.

Good luck!

Name _____

Disc.: Number _____ TA _____ Day/Time _____

“I have adhered to the Duke Community
Standard in completing this
examination.”

1. _____

2. _____

3. _____

4. _____

5. _____

Signature: _____

Total Score _____ (/100 points)

1. (20 pts) In this problem we consider the inner product space $P = \text{span}\{\cos x, \sin x\} \subset C[0, 2\pi]$, using the L^2 inner product.
- (a) Show that $\cos x$ is orthogonal to $\sin x$.
- (b) Find the value k such that $\{k \cos x, k \sin x\}$ is an orthonormal basis for P .
- (c) Suppose that $f = a \cos x + b \sin x$, and that the values of f are known but the values of a and b are not. Find formulas for a and b in terms of f . (Be sure to indicate clearly where you use part (b) in your argument.)

2. (20 pts)

(a) Diagonalize the matrix A below.

$$A = \begin{pmatrix} -1 & 2 \\ -6 & 6 \end{pmatrix}$$

(b) Find a fundamental set of solutions to the system $\vec{y}' = A\vec{y}$.

3. (20 pts)

- (a) The arithmetic below is given. Suppose that B represents a linear transformation T with respect to the standard basis \mathcal{S} . Find the change of basis matrix $[I]_{\mathcal{V}}^{\mathcal{S}}$, where \mathcal{V} is a Jordan basis for T . (Be sure to clearly show your reasoning.)

$$I = \begin{pmatrix} 1 & -4 & 2 \\ 1 & -1 & 0 \\ -1 & 3 & -1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 1 & 3 \end{pmatrix}$$

$$B = \begin{pmatrix} -4 & -3 & -10 \\ -1 & 3 & -1 \\ 5 & 2 & 10 \end{pmatrix} = \begin{pmatrix} 1 & -4 & 2 \\ 1 & -1 & 0 \\ -1 & 3 & -1 \end{pmatrix} \begin{pmatrix} 3 & 1 & 0 \\ 0 & 3 & 1 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 2 \\ 1 & 1 & 2 \\ 2 & 1 & 3 \end{pmatrix}$$

- (b) There is a system $\vec{u}' = J\vec{u}$ (where J is in Jordan form) whose solutions relate to solutions to the system $\vec{y}' = B\vec{y}$. Find this system and the matrix C with $\vec{u} = C\vec{y}$, and show this relationship holds.

4. (20 pts) The only eigenvector for the matrix A below is the indicated vector \vec{v}_1 below.

$$\begin{pmatrix} 5 & -1 \\ 9 & -1 \end{pmatrix} \quad \vec{v}_1 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

Find a fundamental set of solutions to the system $\vec{y}' = A\vec{y}$.

5. (20 pts) Find a pair of matrix equations whose solution (if possible) would result in the vectors \vec{a} and \vec{b} that make $\vec{y}_p = (\vec{a} + \vec{b}x)e^{rx}$ a particular solution to the system

$$\vec{y}' = A\vec{y} + xe^{rx}\vec{v}$$

What is the condition on r that would guarantee that the solution would be possible?