

Homework #4

due Monday, September 20, at 11:59pm

1. For each matrix A and vector b , and express the solution set in the form

$$p + \text{Span}\{???\}$$

for some vector p . For instance,

$$\begin{pmatrix} 1 & -1 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}.$$

[Hint: You found the parametric vector form in Problem 8 of Homework 3.]

a) $A = \begin{pmatrix} 2 & 1 & 1 & 4 \\ 4 & 2 & 1 & 7 \end{pmatrix} \quad b = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

b) $A = \begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix} \quad b = \begin{pmatrix} 3 \\ 2 \\ 49 \end{pmatrix}$

c) $A = \begin{pmatrix} 1 & 2 & 3 & -1 & 1 \\ -2 & -4 & -5 & 4 & 1 \\ 1 & 2 & 2 & -3 & -1 \\ -3 & -6 & -7 & 7 & 6 \end{pmatrix} \quad b = \begin{pmatrix} 2 \\ 4 \\ -6 \\ 10 \end{pmatrix}$

d) $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} \quad b = \begin{pmatrix} 2 \\ 5 \\ 4 \end{pmatrix}$

2. For each matrix A in Problem 1, write the solution set of $Ax = 0$ as a span. Does there exist a nontrivial solution? Do not do Gauss–Jordan elimination again!
3. When is the following system consistent?

$$\begin{aligned} 2x_1 + 2x_2 - x_3 &= b_1 \\ -4x_1 - 5x_2 + 5x_3 &= b_2 \\ 6x_1 + x_2 + 12x_3 &= b_3 \end{aligned}$$

Your answer should be a single linear equation in b_1, b_2, b_3 . Explain the relationship between this equation and

$$\text{Span} \left\{ \begin{pmatrix} 2 \\ -4 \\ 6 \end{pmatrix}, \begin{pmatrix} 2 \\ -5 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 5 \\ 12 \end{pmatrix} \right\}.$$

4. Suppose that A is a 2×3 matrix such that

$$A \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad \text{and} \quad A \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

- a) Find two different solutions of $Ax = 0$.

- b) Find two more solutions of $Ax = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$.
5. Suppose that $Ax = b$ is consistent. Explain why $Ax = b$ has a unique solution precisely when $Ax = 0$ has only the trivial solution.
6. Give geometric descriptions of the following spans (line, plane, ...).

a) $\text{Span} \left\{ \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} \right\}$ b) $\text{Span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 0 \\ -2 \\ 1 \end{pmatrix} \right\}$ c) $\text{Span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 0 \\ -2 \\ -6 \end{pmatrix} \right\}$

d) $\text{Span} \left\{ \begin{pmatrix} 2 \\ -4 \\ 6 \end{pmatrix}, \begin{pmatrix} 2 \\ -5 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 5 \\ 12 \end{pmatrix} \right\}$ e) $\text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} \right\}$

[Hint: for d), compare Problem 3.]

7. a) List five nonzero vectors contained in $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$.
- b) Is $\begin{pmatrix} 0 \\ 3 \\ 6 \end{pmatrix}$ contained in $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$?
- If so, express $\begin{pmatrix} 0 \\ 3 \\ 6 \end{pmatrix}$ as a linear combination of $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}$.
- c) Show that $\begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}$ is contained in $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} \right\}$.
- d) Describe $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$ geometrically.
- e) Find a vector not contained in $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$.
8. Decide if each statement is true or false, and explain why.
- a) A vector b is a linear combination of the columns of A if and only if $Ax = b$ has a solution.
- b) There is a matrix A such that $Ax = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$ has infinitely many solutions and $Ax = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$ has exactly one solution.
- c) The zero vector is contained in every span.

- d) The matrix equation $Ax = 0$ can be consistent or inconsistent, depending on what A is.
- e) If the zero vector is a solution of a system of equations, then the system is homogeneous.
- f) If $Ax = b$ has a unique solution, then A has a pivot in every column.
- g) If $Ax = b$ is consistent, then the solution set of $Ax = b$ is obtained by translating the solution set of $Ax = 0$.
- h) It is possible for $Ax = b$ to have exactly 13 solutions.
9. Find a spanning set for the null space of each matrix, and express the null space as the column space of some other matrix.



$$\text{a) } \begin{pmatrix} 2 & 1 & 1 & 4 \\ 4 & 2 & 1 & 7 \end{pmatrix} \quad \text{b) } \begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix}$$

$$\text{c) } \begin{pmatrix} 1 & 2 & 3 & -1 & 1 \\ -2 & -4 & -5 & 4 & 1 \\ 1 & 2 & 2 & -3 & -1 \\ -3 & -6 & -7 & 7 & 6 \end{pmatrix} \quad \text{d) } \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$$



[Hint: Compare Problem 2.]

10. Draw pictures of the null space and the column space of the following matrices. Be precise!



a) $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$:

	Nul(A)		Col(A)
			

b) $A = \begin{pmatrix} 1 & 2 \\ -1 & -2 \end{pmatrix}$:

	Nul(A)		Col(A)
			

c) $A = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$:

	Nul(A)		Col(A)
			

11. Give examples of subsets V of \mathbf{R}^2 such that:
- V is closed under addition and contains 0, but is not closed under scalar multiplication.
 - V is closed under scalar multiplication and contains 0, but is not closed under addition.
 - V is closed under addition and scalar multiplication, but does not contain 0.
- Therefore, none of these conditions is redundant.
12. Which of the following subsets of \mathbf{R}^3 are subspaces? If it is not a subspace, why not? If it is, write it as the column space or null space of some matrix.
- The plane $\{(x, y, x) : x, y \in \mathbf{R}\}$.
 - The plane $\{(x, y, 1) : x, y \in \mathbf{R}\}$.
 - The set consisting of all vectors (x, y, z) such that $xy = 0$.
 - The set consisting of all vectors (x, y, z) such that $x \leq y$.

e) The span of $(1, 2, 3)$ and $(2, 1, -3)$.

f) The solution set of the system of equations $\begin{cases} x + y + z = 0 \\ x - 2y - z = 0. \end{cases}$

g) The solution set of the system of equations $\begin{cases} x + y + z = 0 \\ x - 2y - z = 1. \end{cases}$

13. Give a geometric description of the following column spaces (line, plane, ...).

a) $\text{Col} \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$ b) $\text{Col} \begin{pmatrix} 0 & 0 \\ 1 & -2 \\ 3 & 1 \end{pmatrix}$ c) $\text{Col} \begin{pmatrix} 0 & 0 \\ 1 & -2 \\ 3 & -6 \end{pmatrix}$

d) $\text{Col} \begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix}$ e) $\text{Col} \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$

[Hint: Compare Problem 6.]

14. Find a nonzero 2×2 matrix such that $A^2 = 0$.

15. a) Explain why $\text{Col}(AB)$ is contained in $\text{Col}(A)$.

b) Give an example where $\text{Col}(AB) \neq \text{Col}(A)$. Can you take $A = B$?
[Hint: use Problem 14.]

16. a) Explain why $\text{Nul}(AB)$ contains $\text{Nul}(B)$.

b) Give an example where $\text{Nul}(AB) \neq \text{Nul}(B)$. Can you take $A = B$?
[Hint: use Problem 14.]

17. a) If $\text{Col}(B)$ is contained in $\text{Nul}(A)$, then $AB = \underline{\hspace{2cm}}$.

b) Find a 2×2 matrix A such that $\text{Col}(A) = \text{Nul}(A)$. What is the rank of such a matrix? [Hint: use Problem 14.]

18. Find a matrix A such that

$$\text{Col}(A) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \right\} \quad \text{and} \quad \text{Nul}(A) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

What is the rank of A ?

19. For the following matrix A , compute the reduced row echelon form of A and of A^T . Do they have the same free variables? Do they have the same rank?

$$A = \begin{pmatrix} 1 & 2 & 3 \\ -1 & -2 & -3 \\ 4 & 5 & 6 \end{pmatrix}$$

- 20.** Decide if each statement is true or false, and explain why.
- a) The null space of an $m \times n$ matrix with n pivots is \mathbf{R}^n .
 - b) If $\text{Col}(A) = \{0\}$, then A is the zero matrix.
 - c) The column space of $2A$ equals the column space of A .
 - d) The null space of $A + B$ contains the null space of A .
 - e) If U is an echelon form of A , then $\text{Nul}(U) = \text{Nul}(A)$.
 - f) If U is an echelon form of A , then $\text{Col}(U) = \text{Col}(A)$.