

## Math 218D Problem Session

Week 11

### 1. Matrices with complex eigenvalues

Consider the matrices  $A = \begin{pmatrix} 0 & -1/2 \\ 1/2 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ .

- a) Compute the eigenvalues of  $A$  and  $B$ . Write each eigenvalue in polar coordinates  $z = re^{i\theta}$ .
- b) Compute the eigenvectors of  $A$  and  $B$ .

## 2. The dynamics of a diagonal matrix

Consider the matrix  $A = \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix}$ .

- a) For each of the following vectors, plot  $v$ ,  $Av$ , and  $A^2v$ :
- (1)  $v = (1, 0)$
  - (2)  $v = (0, 1)$
  - (3)  $v = (1, 1)$
- b) For each of the same vectors, sketch the shape you get by connecting the dots between the points  $\dots, A^{-2}v, A^{-1}v, v, Av, A^2v, \dots$
- c) For the vector  $v = (1, 1)$ , what direction is the vector  $A^n v$  approximately pointing when  $n$  is very large? In other words, what unit vector does  $\frac{A^n v}{\|A^n v\|}$  approximate when  $n$  is very large?
- d) For the vector  $v = (1, 1)$ , what direction is  $A^{-n}v$  approximately pointing when  $n$  is very large?

### 3. The dynamics of a diagonalizable matrix

Consider the matrix  $A$  with  $A(1, 1) = 3(1, 1)$  and  $A(1, -2) = 2(1, -2)$ . In other words,  $A$  is diagonalizable and you have been told the eigenvectors and eigenvalues.

a) For each of the following vectors, plot  $v$ ,  $Av$ ,  $A^2v$ :

(1)  $v = (1, 1)$

(2)  $v = (1, -2)$

(3)  $v = (2, -1)$

**You can do this without computing the matrix  $A$ !**

b) For each of the same vectors, sketch the shape you get by connecting the dots between the points  $\dots, A^{-2}v, A^{-1}v, v, Av, A^2v, \dots$

c) For the vector  $v = (2, -1)$ , what direction is the vector  $A^n v$  approximately pointing when  $n$  is very large?

d) For the vector  $v = (2, -1)$ , what direction is  $A^{-n} v$  approximately pointing when  $n$  is very large?

#### 4. Dynamics of complex matrices

Consider the matrices  $A = \begin{pmatrix} 0 & -1/2 \\ 1/2 & 0 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$  you studied in problem 1.

- a) Plot the points  $(4, 0)$ ,  $A(4, 0)$ ,  $A^2(4, 0)$ ,  $A^3(4, 0)$ , and  $A^4(4, 0)$ . Connect the dots between these points. Predict the shape that you would get if you continued to  $A^5(4, 0)$ ,  $A^6(4, 0)$ ,  $\dots$
- b) Plot the points  $(1, 0)$ ,  $B(1, 0)$ ,  $B^2(1, 0)$ ,  $B^3(1, 0)$ , and  $B^4(1, 0)$ . Connect the dots between these points. Predict the shape that you would get if you continued to  $B^5(1, 0)$ ,  $B^6(1, 0)$ ,  $\dots$
- c) What do the eigenvalues you found in problem 1a) explain about your pictures from a) and b)?
- d) Find complex scalars  $a$ ,  $b$  such that  $(1, 0) = av_1 + bv_2$ , where  $v_1$  and  $v_2$  are the eigenvectors for  $B$  you found in problem 1c).
- e) Compute  $B^n(1, 0)$  in terms of complex exponentials.
- f) Use Euler's formula  $e^{i\theta} = \cos(\theta) + i \sin(\theta)$  to write  $B^n(1, 0)$  in terms of trig. functions (no complex numbers should appear in your final answer).
- g) Can you predict a formula for  $A^n(4, 0)$  in terms of trig. functions?