

## Math 218D Problem Session

Week 1

### 1. Row Echelon Form

a)

$$\left( \begin{array}{cc|c} 1 & 2 & 1 \\ 0 & 3 & 2 \end{array} \right).$$

In REF, pivots are 1 and 3

b)

$$\begin{aligned} 5x - 2y &= 1 \\ 0x + 0y &= 6 \\ 0x + 0y &= 0 \end{aligned}$$

In REF, pivots are 5 and 6

c)

$$\left( \begin{array}{ccc|c} 2 & 1 & -1 & 1 \\ 0 & 3 & 1 & 2 \\ 2 & 0 & 0 & 1 \end{array} \right).$$

Not in REF

d)

$$\begin{aligned} 2x + y &= 3 \\ y &= 5 \end{aligned}$$

In REF, pivots are 2 and 5

e)

$$\begin{aligned} 3x + 2y + z &= 0 \\ 0x + 0y + 0z &= 0 \\ -x - 2y + 4z &= 1 \end{aligned}$$

Not in REF


f)

$$\begin{aligned} 5x + 5y &= 5 \\ x + y &= 1 \end{aligned}$$

Not in REF

### 2. Two Equations and Two Unknowns

a)



b) The linear system is

$$\begin{aligned}x - y &= 2 \\2x - 4y &= -4.\end{aligned}$$

Subtract  $2 \cdot R_1$  from  $R_2$  to obtain:

$$\begin{aligned}x - y &= 2 \\-2y &= -8.\end{aligned}$$

c) Divide the second row by 2 to obtain:

$$\begin{aligned}x - y &= 2 \\y &= 4.\end{aligned}$$

d) Add the second row to the first row to obtain:

$$\begin{aligned}x &= 6 \\y &= 4.\end{aligned}$$

This is the solution.

e)  $6 - 4 = 2$ ,  $2 \cdot 6 - 4 \cdot 4 = -5$ .

f) The system first becomes in REF after the 1st row operation. The pivots are 1 and  $-2$ .

### 3. Three Equations Three Unknowns

a)  $A = \begin{pmatrix} 1 & -3 & 1 \\ 2 & -8 & 8 \\ -6 & 3 & -15 \end{pmatrix}, b = \begin{pmatrix} 4 \\ -2 \\ 9 \end{pmatrix}.$

b) The augmented matrix is

$$\left( \begin{array}{ccc|c} 1 & -3 & 1 & 4 \\ 2 & -8 & 8 & -2 \\ -6 & 3 & -15 & 9 \end{array} \right).$$

c) First, replace  $R_2$  by  $R_2 - 2R_1$  ( $R_2 += -2R_1$ ).

$$\left( \begin{array}{ccc|c} 1 & -3 & 1 & 4 \\ 0 & -2 & 6 & -10 \\ -6 & 3 & -15 & 9 \end{array} \right).$$

Then  $R_3 += 6R_1$ :

$$\left( \begin{array}{ccc|c} 1 & -3 & 1 & 4 \\ 0 & -2 & 6 & -10 \\ 0 & -15 & -9 & 33 \end{array} \right)..$$

Now, you can do row scaling here, although you don't need to. Let's do it now to simplify our rows:  $R_2 \times = -(1/2)$  and  $R_3 \times = -(1/3)$  (combining two elementary operations at once):

$$\left( \begin{array}{ccc|c} 1 & -3 & 1 & 4 \\ 0 & 1 & -3 & 5 \\ 0 & 5 & 3 & -11 \end{array} \right).$$

We do one more row addition, replacing  $R_2$  with  $R_2 - 5R_1$  ( $R_2 -= 5R_1$ ):

$$\left( \begin{array}{ccc|c} 1 & -3 & 1 & 4 \\ 0 & 1 & -3 & 5 \\ 0 & 0 & 18 & -36 \end{array} \right).$$

Do one more row scaling, replacing  $R_3$  with  $\frac{1}{18}R_3$  ( $R_3 \times = 1/18$ ):

$$\left( \begin{array}{ccc|c} 1 & -3 & 1 & 4 \\ 0 & 1 & -3 & 5 \\ 0 & 0 & 1 & -2 \end{array} \right).$$

d) I used 6 elementary row operations, but the row scalings could have been avoided, giving you as few as 3.

e) The system of equations is now

$$\begin{aligned} x_1 - 3x_2 + x_3 &= 4 \\ x_2 - 3x_3 &= 5 \\ x_3 &= -2 \end{aligned}.$$

Substituting  $x_3 = -2$ , we obtain the system

$$\begin{aligned}x_1 - 3x_2 &= 6 \\x_2 &= -1 \\x_3 &= -2\end{aligned}$$

Substituting  $x_2 = -1$ , we obtain the system

$$\begin{aligned}x_1 &= 3 \\x_2 &= -1 \\x_3 &= -2\end{aligned}$$

which is the solution.

f) Check  $\begin{pmatrix} 1 & -3 & 1 \\ 2 & -8 & 8 \\ -6 & 3 & -15 \end{pmatrix} \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 4 \\ -2 \\ 9 \end{pmatrix}$ .

#### 4. Another One—What's Different?

Consider the system of three linear equations

$$\begin{aligned}x_1 - 2x_2 + x_3 &= -2 \\2x_1 - 4x_2 + 8x_3 &= 2 \\x_1 - 3x_2 - x_3 &= 1.\end{aligned}$$

a) The linear system is

$$\begin{aligned}x_1 - 2x_2 + x_3 &= -2 \\2x_1 - 4x_2 + 8x_3 &= 2 \\x_1 - 3x_2 - x_3 &= 1.\end{aligned}$$

By doing two row subtraction operations ( $R_2 - = 2R_1$  and  $R_3 - = R_1$ ), we obtain

$$\begin{aligned}x_1 - 2x_2 + x_3 &= -2 \\6x_3 &= 6 \\-x_2 - 2x_3 &= 3.\end{aligned}$$

b) We swap rows 1 and 2 to obtain

$$\begin{aligned}x_1 - 2x_2 + x_3 &= -2 \\-x_2 - 2x_3 &= 3 \\6x_3 &= 6.\end{aligned}$$

c) Dividing row 3 by 6 gives  $x_3 = 1$ , which we substitute into the first two equations:

$$\begin{aligned}x_1 - 2x_2 &= -3 \\-x_2 &= 5 \\x_3 &= 1.\end{aligned}$$

Dividing row 2 by  $-1$  gives  $x_2 = -5$ , which we substitute into the 1st equation:

$$\begin{aligned}x_1 &= -13 \\x_2 &= -5 \\x_3 &= 1.\end{aligned}$$

This is the solution.

## 5. Traffic Jam

a) We start with

$$120 + w = 250 + x$$

$$120 + x = 70 + y$$

$$390 + y = 250 + z$$

$$115 + z = 175 + w$$

or

$$x - w = -130$$

$$-x + y = 50$$

$$-y + z = 140$$

$$-z + w = -60.$$

b) Eliminating  $x$  from the second equation gives

$$x - w = -130$$

$$y - w = -80$$

$$-y + z = 140$$

$$-z + w = -60.$$

c) Eliminating  $y$  from the third equation gives

$$x - w = -130$$

$$y - w = -80$$

$$z - w = 60$$

$$-z + w = -60.$$

d) Eliminating  $z$  from the fourth equation gives

$$x - w = -130$$

$$y - w = -80$$

$$z - w = 60$$

$$0 + 0 = 0.$$

e) We can't just use substitution, as our final equation is not of the form  $w = (?)$ . The number of cars on roads  $x$ ,  $y$ , and  $z$  all depend on how many cars are on  $w$ .

f) The system has infinitely many solutions. There can be as many cars as you want, travelling in a circle around the town.

g) The augmented matrix is

$$\left( \begin{array}{cccc|c} 1 & 0 & 0 & -1 & -130 \\ 0 & 1 & 0 & -1 & -80 \\ 0 & 0 & 1 & -1 & 60 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right).$$

The pivots are the 1's. Not every row has a pivot. The fourth column does not have a pivot - as we will discuss in week 3, this means that we can find a solution which makes the fourth variable take any value we want. Such a variable is called a *free variable*.