

MATH 218D-1
PRACTICE FINAL EXAMINATION

Name		Duke NetID	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 180 minutes to complete this exam.
- If you finish early, go back and check your work.
- The graders will only see the work on the **printed pages** (front and back). You may use other scratch paper, but the graders will not see anything written there.
- You may use a **four-function calculator** for doing arithmetic, but you should not need one. You may bring a 8.5×11 -**inch note sheet** covered with anything you want. All other materials and aids are strictly prohibited.
- For full credit you must **show your work** so that your reasoning is clear, unless otherwise indicated.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 180 minutes, without notes or distractions.

[This page intentionally left blank]

Problem 1.

[30 points]

Consider the following matrix A and its singular value decomposition:

$$A = \begin{pmatrix} 0 & 6 & -3 \\ 5 & 8 & 1 \\ 1 & -8 & 5 \\ 8 & -4 & 10 \end{pmatrix} = \sigma_1 u_1 v_1^T + \sigma_2 u_2 v_2^T$$

$$\sigma_1 = 3\sqrt{30} \quad u_1 = \frac{1}{\sqrt{30}} \begin{pmatrix} -2 \\ -1 \\ 3 \\ 4 \end{pmatrix} \quad v_1 = \frac{1}{3} \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}$$

$$\sigma_2 = 3\sqrt{15} \quad u_2 = \frac{1}{\sqrt{15}} \begin{pmatrix} 1 \\ 3 \\ -1 \\ 2 \end{pmatrix} \quad v_2 = \frac{1}{3} \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$$

a) The rank of A is $r = \boxed{}$.

b) An orthonormal basis for $\text{Col}(A)$ is:

$$\left\{ \begin{pmatrix} \\ \\ \\ \end{pmatrix} \right\}$$

c) An orthonormal basis for $\text{Row}(A)$ is:

$$\left\{ \begin{pmatrix} \\ \\ \end{pmatrix} \right\}$$

d) An orthonormal basis for $\text{Nul}(A)$ is: (hint: cross products)

$$\left\{ \begin{pmatrix} \\ \\ \\ \end{pmatrix} \right\}$$

e) The *maximum* value of $\|Ax\|$ subject to $\|x\| = 1$ is $\boxed{}$.

f) The *minimum* value of $\|Ax\|$ subject to $\|x\| = 1$ is $\boxed{}$.

[Scratch work for Problem 1]

Problem 1, continued

g) Find a left inverse of A (a matrix B such that $BA = I_3$), or explain why no such matrix exists:

$$B = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

h) The matrix $A^T A$ has an *orthogonal* diagonalization $A^T A = QDQ^T$ for

$$Q = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix} \quad D = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

[Scratch work for Problem 1]

Problem 2.

[25 points]

Consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 2 & 1 \\ 2 & 1 & 1 & 2 \end{pmatrix}.$$

a) Find the singular value decomposition of A in matrix form: $A = U\Sigma V^T$ for

$$U = \begin{pmatrix} & & & \\ & & & \\ & & & \\ & & & \end{pmatrix} \quad \Sigma = \begin{pmatrix} & & & \\ & & & \\ & & & \\ & & & \end{pmatrix} \quad V = \begin{pmatrix} & & & \\ & & & \\ & & & \\ & & & \end{pmatrix}$$

[Scratch work for Problem 2]

[Scratch work for Problem 2]

Problem 2, continued

- e) Compute the orthogonal decomposition of $b = (2, 1, 4, 3)$ with respect to $V_1 = \text{Row}(A)$: $b = b_{V_1} + b_{V_1^\perp}$ for

$$b_{V_1} = \begin{pmatrix} \\ \\ \\ \end{pmatrix} \quad b_{V_1^\perp} = \begin{pmatrix} \\ \\ \\ \end{pmatrix}$$

- f) Compute the orthogonal decomposition of $c = (2, 4)$ with respect to $V_3 = \text{Col}(A)$:
 $c = c_{V_3} + c_{V_3^\perp}$ for

$$c_{V_3} = \begin{pmatrix} \\ \end{pmatrix} \quad c_{V_3^\perp} = \begin{pmatrix} \\ \end{pmatrix}$$

[Scratch work for Problem 2]

Problem 3.

[20 points]

Consider the positive-definite quadratic form

$$q(x_1, x_2, x_3) = 5x_1^2 + 4x_2^2 + 3x_3^2 + 4x_1x_2 + 4x_2x_3.$$

The minimum value of $\|u\|^2$ subject to $q(u) = 1$ is $\|u\|^2 = 1/7$, and the maximum value of $\|u\|^2$ subject to $q(u) = 1$ is $\|u\|^2 = 1$

a) Find a vector u_1 satisfying $\|u_1\|^2 = 1/7$ and $q(u_1) = 1$.

$$u_1 = \begin{pmatrix} \\ \\ \end{pmatrix}$$

b) Find a vector u_3 satisfying $\|u_3\|^2 = 1$ and $q(u_3) = 1$.

$$u_3 = \begin{pmatrix} \\ \\ \end{pmatrix}$$

c) The maximum value of $\|u\|^2$ subject to $q(u) = 1$ and $u \perp u_1$ is $\|u\|^2 = 1/4$. Find a vector u_2 satisfying $\|u_2\|^2 = 1/4$ and $q(u_2) = 1$ and $u_2 \perp u_1$.

$$u_2 = \begin{pmatrix} \\ \\ \end{pmatrix}$$

d) Find a change of coordinates $x = Qy$ such that $q(y)$ is diagonal:

$$x_1 =$$

$$x_2 =$$

$$x_3 =$$

[Scratch work for Problem 3]

Problem 4.

[20 points]

Consider the matrix $A = \begin{pmatrix} 1 & 2 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$.

a) Find the QR decomposition of A .

$$Q = \begin{pmatrix} & \\ & \\ & \end{pmatrix} \quad R = \begin{pmatrix} & \\ & \end{pmatrix}$$

b) Find the least squares solution of $Ax = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ using your answer to a).

$$\hat{x} = \begin{pmatrix} \\ \end{pmatrix}$$

c) Find the orthogonal projection b_V of $b = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ onto $V = \text{Col}(A)$.

$$b_V = \begin{pmatrix} \\ \end{pmatrix}$$

d) The singular value decomposition of R is $R = U\Sigma V^T$ for:

$$U = \frac{1}{\sqrt{5}} \begin{pmatrix} \sqrt{3} & -\sqrt{2} \\ \sqrt{2} & \sqrt{3} \end{pmatrix} \quad \Sigma = \begin{pmatrix} \sqrt{6} & 0 \\ 0 & 1 \end{pmatrix} \quad V = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 & -2 \\ 2 & 1 \end{pmatrix}.$$

Find the right singular vectors of A without doing any calculations.

$$v_1 = \begin{pmatrix} \\ \end{pmatrix} \quad v_2 = \begin{pmatrix} \\ \end{pmatrix}$$

[Scratch work for Problem 4]

Problem 5.

[15 points]

Consider the matrix

$$A_0 = \begin{pmatrix} 1 & 3 & 2 & 4 & 5 \\ 8 & 2 & 3 & 9 & 8 \\ 3 & 1 & 4 & 5 & 2 \end{pmatrix}$$

whose columns contain five data points of three measurements each.

a) The averages of the rows are

$$\mu_1 = \boxed{} \quad \mu_2 = \boxed{} \quad \mu_3 = \boxed{}$$

and the centered data matrix is

$$A = \begin{pmatrix} & & & & \\ & & & & \\ & & & & \end{pmatrix}$$

b) The covariance matrix is

$$S = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

The first measurement has variance $s_1^2 = \boxed{}$ and the total variance is $s^2 = \boxed{}$.

The eigenvalues and unit eigenvectors of S are

$$\begin{aligned} \lambda_1 &\approx 11.3 & \lambda_2 &\approx 2.74 & \lambda_3 &\approx 1.45 \\ u_1 &\approx \begin{pmatrix} 0.156 \\ 0.958 \\ 0.240 \end{pmatrix} & u_2 &\approx \begin{pmatrix} -0.792 \\ -0.024 \\ 0.610 \end{pmatrix} & u_3 &\approx \begin{pmatrix} -0.590 \\ 0.286 \\ -0.755 \end{pmatrix}. \end{aligned}$$

c) In the sense of orthogonal least squares, the best-fit...

$$\text{line is } L = \text{Span} \left\{ \begin{pmatrix} \\ \\ \end{pmatrix} \right\} \quad \text{and plane is } V = \text{Span} \left\{ \begin{pmatrix} \\ \\ \end{pmatrix}, \begin{pmatrix} \\ \\ \end{pmatrix} \right\}.$$

The variance along L is $\boxed{}$ and the variance along V is $\boxed{}$.

[Scratch work for Problem 5]

Problem 6.

[25 points]

Fill in the circles of all choices that apply. No justification is necessary.

a) Let A be a 3×3 matrix such that $Ax = (2, 1, 0)$ does not have a solution. Which of the following are *impossible*?

- A is invertible.
- $Ax = (1, 2, 0)$ does not have a solution.
- The solution set of $Ax = (1, 2, 0)$ is $\{(1, 1, 0)\}$.
- The solution set of $Ax = (4, 2, 0)$ is a line.
- The solution set of $Ax = (2, 4, 0)$ is a line through the origin.

b) Which of the following sets form a basis for $\text{Nul}\left(\begin{smallmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{smallmatrix}\right)$?

- $\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \end{pmatrix} \right\}$
- $\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} \right\}$
- $\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix} \right\}$
- $\left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} \right\}$
- $\left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} -2 \\ 2 \\ -2 \\ 2 \end{pmatrix} \right\}$
- $\left\{ \begin{pmatrix} 4 \\ -4 \\ 4 \\ -4 \end{pmatrix} \right\}$

c) Which of the following quadratic forms are positive-definite?

- $q(x_1, x_2, x_3) = x_1^2 - 2x_2^2 + 3x_3^2 + 4x_1x_2 + 2x_1x_3$
- $q(x_1, x_2, x_3) = 11x_1^2 + 10x_2^2 + 13x_3^2 + 12x_1x_2 + 6x_1x_3 + 2x_2x_3$
- $q(x_1, x_2, x_3) = (x_1 + x_2 - x_3)^2$
- $q(x) = x^T \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} x$

d) Which of the following matrices are diagonalizable over the real numbers?

- $\begin{pmatrix} 1 & 7 & 5 \\ 7 & 2 & 4 \\ 5 & 4 & 9 \end{pmatrix}$
- $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
- $\begin{pmatrix} 2 & 7 & 1 & 4 \\ 0 & 1 & 3 & 9 \\ 0 & 0 & -1 & -12 \\ 0 & 0 & 0 & 3 \end{pmatrix}$
- $\begin{pmatrix} \cos(27^\circ) & -\sin(27^\circ) \\ \sin(27^\circ) & \cos(27^\circ) \end{pmatrix}$
- $\left(\begin{array}{l} \text{The matrix} \\ \text{for projection} \\ \text{onto} \end{array} \right) \text{Span} \left\{ \begin{pmatrix} 1 \\ 3 \\ 2 \\ 7 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \\ 2 \\ 4 \end{pmatrix} \right\}$

e) Which of the following phrases do *not* make sense? The letter A refers to a matrix.

- The dimension of A is equal to 5.
- A is linearly independent.
- The subspace of A has dimension 4.
- The solution set of $Ax = (1, 2)$ is empty.
- The orthogonal complement of A is a plane.

[Scratch work for Problem 6]

[Scratch work for Problem 7]

Problem 8.

[30 points]

True/false problems: **circle** the correct answer. No justification is needed. *All matrices in this problem have real entries.*

- a) **T** **F** The singular values of a symmetric matrix are the absolute values of the eigenvalues.
- b) **T** **F** If A has linearly independent columns, then A^+A is the identity matrix.
- c) **T** **F** A positive-definite matrix has positive numbers on the main diagonal.
- d) **T** **F** A diagonalizable 5×5 matrix has 5 different eigenvalues.
- e) **T** **F** Every subspace of \mathbf{R}^n has an orthonormal basis.
- f) **T** **F** If $V = \{(x, y, z) \in \mathbf{R}^3 : x = 2y\}$, then V^\perp is a line.
- g) **T** **F** If V is a subspace and x is not in V , then the orthogonal projection of x onto V is zero.
- h) **T** **F** If V is a subspace and x is not in V , then the orthogonal projection of x onto V^\perp is nonzero.
- i) **T** **F** If $\{v_1, \dots, v_n\}$ is a basis for \mathbf{R}^5 , then $n = 5$.
- j) **T** **F** If λ is an eigenvalue of AA^T then λ is an eigenvalue of $A^T A$.

[Scratch work for Problem 8]

Problem 9.

[15 points]

Consider the positive-definite quadratic form

$$q(x_1, x_2) = 4x_1^2 + 4x_2^2 + 2x_1x_2.$$


- a) Find a change of variables $x = Qy$ such that $q(y)$ is diagonal, and write the diagonal form $q(y)$:

$$x_1 =$$

$$x_2 =$$

$$q(y_1, y_2) =$$

- b) Draw the set of all points (x_1, x_2) satisfying $q(x_1, x_2) = 1$ on the grid below. Be precise!



grid lines are 0.1 unit apart

[Scratch work for Problem 9]

Problem 10.

[15 points]


A centered data matrix A has 2 rows and 10 columns. Its SVD has the form

$$A = 5u_1v_1^T + 2u_2v_2^T,$$

where u_1, u_2 and v_1, v_2 are the singular vectors. The columns of A and the first left singular vector u_1 are drawn below. *Draw and label:*

- the best-fit line in the sense of orthogonal least squares;
- the direction of *smallest* variance;
- the columns of $5u_1v_1^T$ (drawn as dots).

The variance in the direction of *smallest* variance is .



[Scratch work for Problem 10]