

### Math 218D-1: Homework #3


due Wednesday, September 21, at 11:59pm

1. Consider the vectors

$$v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \quad w = \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

Draw the 16 linear combinations  $cv + dw$  ( $c, d = -1, 0, 1, 2$ ) as *points* in the  $xy$ -plane.

2. Certain vectors  $v, w$  in  $\mathbf{R}^2$  are drawn below. Express each of  $b_1, b_2, b_3, b_4, b_5$  as a linear combination of  $v, w$ . *Do not try to guess the coordinates of  $v$  and  $w$ !*



3. If

$$v + w = \begin{pmatrix} -4 \\ 1 \end{pmatrix} \quad \text{and} \quad v - w = \begin{pmatrix} 2 \\ 3 \end{pmatrix},$$

compute and draw the vectors  $v$  and  $w$ .

4. Consider the vectors

$$u = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad v = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Draw a picture of all of the linear combinations  $au + bv$  for real numbers  $a, b$  satisfying  $0 \leq a \leq 1$  and  $0 \leq b \leq 1$ . (This will be a shaded region in the  $xy$ -plane.)

5. For each matrix  $A$  and vector  $b$ , decide if the system  $Ax = b$  is consistent. If so, find the parametric form of the general solution of  $Ax = b$ . For instance,

$$\begin{pmatrix} 1 & -1 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \rightsquigarrow x_1 = x_2 + 1.$$

Also answer the following questions: Which variables are free? How many solutions does the system have?

a)  $A = \begin{pmatrix} 2 & 1 & 1 & 4 \\ 4 & 2 & 1 & 7 \end{pmatrix} \quad b = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

b)  $A = \begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix} \quad b = \begin{pmatrix} 3 \\ 2 \\ 49 \end{pmatrix}$

c)  $A = \begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix} \quad b = \begin{pmatrix} 3 \\ 2 \\ 48 \end{pmatrix}$

d)  $A = \begin{pmatrix} 1 & 2 & 3 & -1 & 1 \\ -2 & -4 & -5 & 4 & 1 \\ 1 & 2 & 2 & -3 & -1 \\ -3 & -6 & -7 & 7 & 6 \end{pmatrix} \quad b = \begin{pmatrix} 2 \\ 4 \\ -6 \\ 10 \end{pmatrix}$

e)  $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} \quad b = \begin{pmatrix} 2 \\ 5 \\ 4 \end{pmatrix}$

6. For each matrix  $A$  and vector  $b$  in Problem 5, find the parametric vector form of the general solution of  $Ax = b$  (if the system is consistent). For instance,

$$\begin{pmatrix} 1 & -1 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

What is the dimension of the solution set?

7. The equation  $x + 2y = z$  determines a plane in  $\mathbf{R}^3$ . (This is an *implicit equation* for the plane).
- What is the coefficient matrix  $A$  for this system?
  - Which are the free variables?
  - Write the parametric form of the solutions of  $x + 2y = z$ . This expresses the points on the plane in terms of two *parameters*.
  - Do the same for the plane defined by  $2y = z$ . What is different?

8. The equations

$$\begin{aligned}x + y + z &= 0 \\x - 2y - z &= 1\end{aligned}$$

determine a line  $\mathbf{R}^3$ . (These are *implicit equations* for the line). Write the line in parameterized form: that is, find three linear functions  $f_1(t), f_2(t), f_3(t)$  in one variable such that all points on the line have the form  $(x, y, z) = (f_1(t), f_2(t), f_3(t))$  for a unique value of  $t$ . (Use the free variable as the parameter  $t$ .)

9. Find a  $2 \times 3$  matrix  $A$  in RREF and a vector  $b$  such that the solution set of  $Ax = b$  consists of all vectors of the form

$$\begin{pmatrix} 1+t \\ 2-t \\ t \end{pmatrix} \quad t \in \mathbf{R}.$$

10. Decide if each statement is true or false, and explain why.

- a) A square matrix has no free variables.
- b) An invertible matrix has no free variables.
- c) An  $m \times n$  matrix has at most  $m$  pivots.
- d) A wide matrix (more columns than rows) must have a free variable.
- e) If  $A$  is a tall matrix (more rows than columns), then  $Ax = b$  has at most one solution.

11. Express each system of linear equations as a vector equation. For example,

$$\begin{aligned}x_1 + 2x_2 &= 3 \\ -x_1 - x_2 &= 4\end{aligned} \quad \rightsquigarrow \quad x_1 \begin{pmatrix} 1 \\ -1 \end{pmatrix} + x_2 \begin{pmatrix} 2 \\ -1 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \end{pmatrix}.$$

$$\text{a) } \begin{cases} 3x_1 + 2x_2 + 4x_3 = 9 \\ -x_1 + 4x_3 = 2 \end{cases} \quad \text{b) } \begin{pmatrix} 3 & -5 \\ 2 & 4 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$$

$$\text{c) } \left( \begin{array}{cccc|c} 1 & 0 & 1 & 1 & 2 \\ 0 & 3 & -1 & -2 & 4 \\ 1 & -3 & -4 & -3 & 2 \\ 6 & 5 & -1 & -8 & 1 \end{array} \right)$$

12. a) Is  $\begin{pmatrix} 3 \\ 2 \\ 49 \end{pmatrix}$  a linear combination of  $\begin{pmatrix} 2 \\ -4 \\ 6 \end{pmatrix}$ ,  $\begin{pmatrix} 2 \\ -5 \\ 1 \end{pmatrix}$ ,  $\begin{pmatrix} -1 \\ 5 \\ 12 \end{pmatrix}$ ?

If so, what are the weights?

b) Find a vector that is *not* a linear combination of the columns of the matrix

$$\begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix}.$$

[Hint: for both parts, compare Problem 5.]

13. For each matrix  $A$  and vector  $b$ , and express the solution set in the form

$$p + \text{Span}\{\text{??}\}$$

for some vector  $p$ . For instance,

$$\begin{pmatrix} 1 & -1 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}.$$

[Hint: You found the parametric vector form in Problem 6.]

a)  $A = \begin{pmatrix} 2 & 1 & 1 & 4 \\ 4 & 2 & 1 & 7 \end{pmatrix} \quad b = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

b)  $A = \begin{pmatrix} 2 & 2 & -1 \\ -4 & -5 & 5 \\ 6 & 1 & 12 \end{pmatrix} \quad b = \begin{pmatrix} 3 \\ 2 \\ 49 \end{pmatrix}$

c)  $A = \begin{pmatrix} 1 & 2 & 3 & -1 & 1 \\ -2 & -4 & -5 & 4 & 1 \\ 1 & 2 & 2 & -3 & -1 \\ -3 & -6 & -7 & 7 & 6 \end{pmatrix} \quad b = \begin{pmatrix} 2 \\ 4 \\ -6 \\ 10 \end{pmatrix}$

d)  $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} \quad b = \begin{pmatrix} 2 \\ 5 \\ 4 \end{pmatrix}$

14. For each matrix  $A$  in Problem 13, write the solution set of  $Ax = 0$  as a span. Does there exist a nontrivial solution? Do not do Gauss–Jordan elimination again!

15. Suppose that  $A$  is a  $3 \times 3$  matrix and  $b$  is a vector such that the solution set of  $Ax = b$  is a line in  $\mathbf{R}^3$ . How many pivots does  $A$  have?

16. When is the following system consistent?

$$\begin{aligned} 2x_1 + 2x_2 - x_3 &= b_1 \\ -4x_1 - 5x_2 + 5x_3 &= b_2 \\ 6x_1 + x_2 + 12x_3 &= b_3 \end{aligned}$$

Your answer should be a single linear equation in  $b_1, b_2, b_3$ . [Hint: perform Gaussian elimination.] Explain the relationship between your answer and

$$\text{Span} \left\{ \begin{pmatrix} 2 \\ -4 \\ 6 \end{pmatrix}, \begin{pmatrix} 2 \\ -5 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 5 \\ 12 \end{pmatrix} \right\}.$$

17. Let  $A$  be a  $3 \times 4$  matrix whose columns span the plane  $x + y + z = 0$ .
- Find a vector  $b \in \mathbf{R}^3$  making the system  $Ax = b$  consistent.
  - Find a vector  $b \in \mathbf{R}^3$  making the system  $Ax = b$  inconsistent.

18. Draw a picture of all vectors  $b \in \mathbf{R}^2$  for which the equation

$$\begin{pmatrix} 1 & 2 \\ -2 & -4 \end{pmatrix} x = b$$

is consistent. [Hint: the answer is a span!]

19. Suppose that  $A$  is a  $2 \times 3$  matrix such that

$$A \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad \text{and} \quad A \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

- Find two different solutions of  $Ax = 0$ .
  - Find two more solutions of  $Ax = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ .
20. Suppose that  $Ax = b$  is consistent. Explain why  $Ax = b$  has a unique solution precisely when  $Ax = 0$  has only the trivial solution.
21. Give geometric descriptions of the following spans (line, plane, ...).

a)  $\text{Span} \left\{ \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} \right\}$       b)  $\text{Span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 0 \\ -2 \\ 1 \end{pmatrix} \right\}$       c)  $\text{Span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 0 \\ -2 \\ -6 \end{pmatrix} \right\}$

d)  $\text{Span} \left\{ \begin{pmatrix} 2 \\ -4 \\ 6 \end{pmatrix}, \begin{pmatrix} 2 \\ -5 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 5 \\ 12 \end{pmatrix} \right\}$       e)  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} \right\}$

[Hint: for d), compare Problem 16.]

22. a) List five nonzero vectors contained in  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$ .

b) Is  $\begin{pmatrix} 0 \\ 3 \\ 6 \end{pmatrix}$  contained in  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$ ?

If so, express  $\begin{pmatrix} 0 \\ 3 \\ 6 \end{pmatrix}$  as a linear combination of  $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}$ .

c) Show that  $\begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}$  is contained in  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} \right\}$ .

d) Describe  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$  geometrically.

e) Find a vector not contained in  $\text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}, \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} \right\}$ .

**23.** Decide if each statement is true or false, and explain why.

- a) A vector  $b$  is a linear combination of the columns of  $A$  if and only if  $Ax = b$  has a solution.
- b) There is a matrix  $A$  such that  $Ax = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$  has infinitely many solutions and  $Ax = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$  has exactly one solution.
- c) The zero vector is contained in every span.
- d) The matrix equation  $Ax = 0$  can be consistent or inconsistent, depending on what  $A$  is.
- e) If the zero vector is a solution of a system of equations, then the system is homogeneous.
- f) If  $Ax = b$  has a unique solution, then  $A$  has a pivot in every column.
- g) If  $Ax = b$  is consistent, then the solution set of  $Ax = b$  is obtained by translating the solution set of  $Ax = 0$ .
- h) It is possible for  $Ax = b$  to have exactly 13 solutions.