

MATH 218D-1
PRACTICE FINAL EXAMINATION

Name		Duke NetID	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 180 minutes to complete this exam.
- If you finish early, go back and check your work.
- The graders will only see the work on the **printed pages** (front and back). You may use other scratch paper, but the graders will not see anything written there.
- You may use a **simple calculator** for doing arithmetic. You may bring a 8.5×11 -inch **note sheet** covered with anything you want. All other materials and aids are strictly prohibited.
- For full credit you must **show your work** so that your reasoning is clear, unless otherwise indicated.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 180 minutes, without notes or distractions.

[This page intentionally left blank]

Problem 1.

[25 points]

Consider the matrix

$$A = \begin{pmatrix} -2/3 & 8/3 & 2/3 \\ & 2 & 2 \\ & & 3 \end{pmatrix}.$$

a) The rank of A is $r = \square$.

b) Explain why it is better to orthogonally diagonalize AA^T instead of $A^T A$ when computing the SVD of A .

Here is the matrix AA^T :

$$AA^T = \begin{pmatrix} 8 & 6 \\ 6 & 17 \end{pmatrix}.$$

c) Compute the characteristic polynomial of AA^T .

$$p(\lambda) =$$

d) Compute the eigenvalues of AA^T .

$$\lambda_1 = \square \quad \lambda_2 = \square$$

e) Compute an orthonormal eigenbasis of AA^T .

$$u_1 = \begin{pmatrix} \\ \end{pmatrix} \quad u_2 = \begin{pmatrix} \\ \end{pmatrix}$$

[Scratch work for Problem 1]

(Problem 1, continued)

f) Compute the SVD of A in outer product form.

$$A = \boxed{} \begin{pmatrix} \\ \end{pmatrix} \begin{pmatrix} \\ \end{pmatrix}^T + \boxed{} \begin{pmatrix} \\ \end{pmatrix} \begin{pmatrix} \\ \end{pmatrix}^T$$

g) Compute the SVD of A in matrix form: $A = U\Sigma V^T$ for

$$U = \begin{pmatrix} \\ \end{pmatrix} \quad \Sigma = \begin{pmatrix} \\ \end{pmatrix} \quad V = \begin{pmatrix} \\ \end{pmatrix}$$

h) What are the eigenvalues of $A^T A$?

[Scratch work for Problem 1]

Problem 2.

[35 points]

Consider the following matrix and its SVD:

$$A = \begin{pmatrix} -2 & 4 \\ 4 & 2 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \\ 1/\sqrt{3} & -1/\sqrt{2} & 1/\sqrt{6} \\ 1/\sqrt{3} & 0 & -2/\sqrt{6} \end{pmatrix} \begin{pmatrix} \sqrt{30} & 0 \\ 0 & 2\sqrt{5} \\ 0 & 0 \end{pmatrix} \frac{1}{\sqrt{10}} \begin{pmatrix} 1 & -3 \\ 3 & 1 \end{pmatrix}^T$$

a) Find an orthonormal basis for $\text{Col}(A)$.

$$\left\{ \begin{pmatrix} \\ \\ \end{pmatrix} \right\}$$

b) Find an orthonormal basis for $\text{Nul}(A^T)$.

$$\left\{ \begin{pmatrix} \\ \\ \end{pmatrix} \right\}$$

c) Write the SVD of A in outer product form.

$$A = \boxed{} \begin{pmatrix} \\ \\ \end{pmatrix} \begin{pmatrix} \\ \end{pmatrix}^T + \boxed{} \begin{pmatrix} \\ \\ \end{pmatrix} \begin{pmatrix} \\ \end{pmatrix}^T$$

[Scratch work for Problem 2]


(Problem 2, continued)

d) Explain why you know $A^T A = \begin{pmatrix} 21 & 3 \\ 3 & 29 \end{pmatrix}$ is positive-definite without doing any calculations.

e) Find an orthogonal diagonalization of $A^T A$: that is, $A^T A = QDQ^T$ for

$$Q = \begin{pmatrix} & \\ & \end{pmatrix}$$
$$D = \begin{pmatrix} & \\ & \end{pmatrix}$$

f) Draw the ellipse defined by $q(x_1, x_2) = 21x_1^2 + 29x_2^2 + 6x_1x_2 = 1$. Grid lines are 0.05 units apart.



g) Find the maximum value of $\|Ax\|^2$ subject to $\|x\| = 1$. At which vectors x is the maximum achieved?

$$\max = \boxed{} \quad x = \begin{pmatrix} \\ \end{pmatrix}$$

h) The minimum value of $\|Ax\|^2$ subject to $\|x\| = 1$ is $\boxed{}$.

[Scratch work for Problem 2]

Problem 3.

[20 points]

Consider the following matrix whose columns consist of 3 data points with 3 measurements each:

$$A_0 = \begin{pmatrix} 4 & 8 & 0 \\ 8 & 1 & 6 \\ 6 & 1 & 2 \end{pmatrix}$$

a) Compute the recentered matrix A (subtract the means of the measurements).

$$A = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

The matrix A has SVD

$$A = 3\sqrt{6} \frac{1}{3} \begin{pmatrix} -2 \\ 2 \\ 1 \end{pmatrix} \frac{1}{\sqrt{6}} (1 \quad -2 \quad 1) + 3\sqrt{2} \frac{1}{3} \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix} \frac{1}{\sqrt{2}} (1 \quad 0 \quad -1)$$

and covariance matrix

$$S = \frac{1}{2} AA^T = \begin{pmatrix} 16 & -10 & -2 \\ -10 & 13 & 8 \\ -2 & 8 & 7 \end{pmatrix}.$$

b) The rank of A is .

c) The total variance is $s^2 =$.

d) The variance of the first measurement is $s_1^2 =$.

e) The variance in the direction $u = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ is $s(u)^2 =$.

f) The variance along the plane $V = \text{Span} \left\{ \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right\}^\perp$ is $s(V)^2 =$.

[Scratch work for Problem 3]

(Problem 3, continued)

g) The line of best fit of the recentered data points is spanned by $u_1 =$.

The variance along this line is $s(u_1)^2 =$, and the error² (the variance along the perpendicular plane) is .

h) The direction of second-largest variance is $u_2 =$, and the variance in this direction is $s(u_2)^2 =$.

i) Find the plane containing the original 3 data points (the columns of A_0) in the form $p + \text{Span}\{\dots\}$.

$$\left(\begin{array}{c} \\ \\ \end{array} \right) + \text{Span} \left\{ \begin{array}{c} \\ \\ \end{array} \right\}$$

[Scratch work for Problem 3]

Problem 4.

[10 points]

a) Compute the characteristic polynomial of the matrix

$$A_1 = \begin{pmatrix} 1 & 3 & 2 \\ 2 & 2 & 1 \\ 1 & 0 & 1 \end{pmatrix}.$$

$$p(\lambda) =$$

b) The matrix

$$A_2 = \begin{pmatrix} 2 & 6 & 6 \\ 3 & -13 & -30 \\ -1 & 6 & 13 \end{pmatrix}$$

has characteristic polynomial

$$p(\lambda) = -(\lambda - 1)(\lambda - 2)(\lambda + 1).$$

Find an invertible matrix C and a diagonal matrix D such that $A = CDC^{-1}$.

$$C = \left(\begin{array}{ccc} & & \\ & & \\ & & \end{array} \right)$$
$$D = \left(\begin{array}{ccc} & & \\ & & \\ & & \end{array} \right)$$

[Scratch work for Problem 4]

Problem 5.

[15 points]

The matrix

$$A = \begin{pmatrix} -1/2 & 10/3 & 25/3 \\ 7/2 & -31/3 & -85/3 \\ -3/2 & 14/3 & 38/3 \end{pmatrix}$$

has a factorization $A = CDC^{-1}$ for

$$C = \begin{pmatrix} 0 & 5 & 2 \\ 5 & -1 & -2 \\ -2 & 1 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1/3 \end{pmatrix}.$$

a) Compute C^{-1} .

$$C^{-1} = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

b) If $v_0 = (3, 6, -2)$, compute $v_k = A^k v_0$: $v_k =$

c) $\lim_{k \rightarrow \infty} A^k v_0 = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}.$

[Scratch work for Problem 5]

Problem 6.


[15 points]

A certain 2×5 matrix A has singular value decomposition

$$A = \sigma_1 u_1 v_1^T + \sigma_2 u_2 v_2^T.$$

The columns of A are drawn as dots on the grid below, and the right singular vectors are drawn as arrows.

- Draw the columns of $\sigma_1 u_1 v_1^T$ as dots on the grid.
- Draw the columns of $\sigma_2 u_2 v_2^T$ as arrows on the grid.
- Explain which geometric quantity in the picture corresponds to σ_2^2 .



[Scratch work for Problem 6]

Problem 7.


[10 points]

a) Find *all* least-squares solutions of $Ax = b$ in parametric vector form, where

$$A = \begin{pmatrix} 1 & 2 \\ 1 & 2 \\ -1 & -2 \end{pmatrix} \quad b = \begin{pmatrix} 6 \\ 0 \\ 3 \end{pmatrix}.$$

$$x = \begin{pmatrix} \quad \\ \quad \end{pmatrix} + \text{Span} \left\{ \begin{array}{l} \quad \\ \quad \end{array} \right\}$$

b) Draw a picture of your answer to a) in the grid below.



[Scratch work for Problem 7]

Problem 8.

[30 points]

Short-answer questions: no justification is necessary.

- a) Find the LDL^T decomposition of the positive-definite, symmetric matrix

$$A = \begin{pmatrix} 2 & 6 \\ 6 & 21 \end{pmatrix} \rightsquigarrow L = \begin{pmatrix} & \\ & \\ & \end{pmatrix} D = \begin{pmatrix} & \\ & \\ & \end{pmatrix}$$

- b) Consider the plane

$$V = \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right\}.$$

List three different methods you could use to compute the projection matrix P_V .

- c) Suppose that A is not diagonalizable and that $A \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$. What is the characteristic polynomial of A ?

$$p(\lambda) =$$

- d) Which of the following sets form a basis for the plane

$$V = \{(x, y, z) \in \mathbf{R}^3 : 3x + 2y + z = 0\}?$$

Fill in the bubbles of all that apply.

$$\begin{aligned} & \circ \left\{ \begin{pmatrix} -1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \\ -6 \end{pmatrix} \right\} \quad \circ \left\{ \begin{pmatrix} -1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \right\} \quad \circ \left\{ \begin{pmatrix} -1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ -1 \end{pmatrix} \right\} \\ & \circ \left\{ \begin{pmatrix} -1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} \right\} \quad \circ \left\{ \begin{pmatrix} -1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\} \quad \circ \left\{ \begin{pmatrix} -1 \\ 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \right\} \end{aligned}$$

- e) Let A be a 4×4 matrix such that 2 and $1 + i$ are eigenvalues of A . Suppose that the 2-eigenspace of A is a plane. Then $\det(A) = \square$.

- f) Suppose that the solution set of $Ax = \begin{pmatrix} 3 \\ 3 \end{pmatrix}$ is the line $\begin{pmatrix} 1 \\ 2 \end{pmatrix} + \text{Span}\left\{\begin{pmatrix} -1 \\ 1 \end{pmatrix}\right\}$. What is the solution set of $Ax = \begin{pmatrix} -3 \\ -3 \end{pmatrix}$?

[Scratch work for Problem 8]

Problem 9.

[20 points]

True/false problems: **circle** the correct answer. No justification is needed.

All matrices in this problem have real entries.

- a) **T** **F** If V is a subspace of \mathbf{R}^n , then $P_V P_{V^\perp} = I_n$.
- b) **T** **F** If $Ax = b$ has infinitely many solutions for every vector $b \in \mathbf{R}^m$, then A has full row rank.
- c) **T** **F** The maximum value of $\|Ax\|^2$ subject to $\|x\| = 1$ is the same as the maximum value of $\|A^T y\|^2$ subject to $\|y\| = 1$.
- d) **T** **F** If σ is a singular value of an invertible matrix A , then σ^{-1} is a singular value of A^{-1} .
- e) **T** **F** If A has linearly independent columns, then $A^T A$ is positive-semidefinite.
- f) **T** **F** If $\{v_1, v_2, v_3, v_4\}$ is a basis of \mathbf{R}^n , then $n = 4$.
- g) **T** **F** Every elementary matrix is diagonalizable.
- h) **T** **F** A diagonalizable 5×5 matrix has 5 distinct eigenvalues.
- i) **T** **F** If A is an $n \times n$ matrix with linearly dependent rows, then the columns of A do not span \mathbf{R}^n .
- j) **T** **F** The diagonal entries of a square matrix are its eigenvalues.

[Scratch work for Problem 9]