

### Math 218D-1: Homework #10

due Wednesday, March 27, at 11:59pm

1. For each  $2 \times 2$  matrix  $A$ , **i)** compute the characteristic polynomial using the formula  $p(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A)$ . Use this to **ii)** find all real eigenvalues, and **iii)** find a basis for each eigenspace, using HW9#14 when applicable. **iv)** Draw and label each eigenspace. **v)** Is the matrix diagonalizable (over the real numbers)?

a)  $\begin{pmatrix} 1 & -2 \\ 1 & 4 \end{pmatrix}$     b)  $\begin{pmatrix} -1 & 1 \\ -9 & 5 \end{pmatrix}$     c)  $\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$     d)  $\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$     e)  $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$

2. For each matrix  $A$ , **i)** find all real eigenvalues of  $A$ , and **ii)** find a basis for each eigenspace. **iii)** Is the matrix diagonalizable (over the real numbers)?

You will probably want to use a computer algebra system to find the roots of the characteristic polynomial. To do so in Sympy, you would type something like:

```
print(roots(-x**3 + 13/4*x + 3/2, multiple=True))  
# [-1.5, -0.5, 2.0]
```

a)  $\begin{pmatrix} -1 & 7 & 5 \\ 0 & 1 & -2 \\ 0 & 1 & 4 \end{pmatrix}$     b)  $\begin{pmatrix} 7 & 12 & 12 \\ -8 & -13 & -12 \\ 4 & 6 & 5 \end{pmatrix}$     c)  $\begin{pmatrix} 6 & 2 & 3 \\ -14 & -7 & -12 \\ 1 & 2 & 4 \end{pmatrix}$

**Optional** (if you want more practice):

d)  $\begin{pmatrix} -11 & -54 & 10 \\ -2 & -7 & 2 \\ -21 & -90 & 20 \end{pmatrix}$     e)  $\begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$

f)  $\begin{pmatrix} 13 & 18 & -18 \\ -12 & -17 & 18 \\ -4 & -6 & 7 \end{pmatrix}$     g)  $\begin{pmatrix} -10 & 28 & -18 & -76 \\ -1 & 9 & -6 & -2 \\ 4 & -8 & 7 & 26 \\ 0 & 2 & -2 & 4 \end{pmatrix}$

3. Let  $V$  be the plane  $x + y + z = 0$ , and let  $R_V = I_3 - 2P_{V^\perp}$  be the reflection matrix over  $V$ , as in HW9#6. Find an eigenbasis for  $R_V$  without doing any computations. Is  $R_V$  diagonalizable?

4. The *Fibonacci numbers* are defined recursively as follows:

$$F_0 = 0, \quad F_1 = 1, \quad F_{n+2} = F_{n+1} + F_n \quad (n \geq 0).$$

The first few Fibonacci numbers are 0, 1, 1, 2, 3, 5, 8, 13, ... In this problem, you will find a closed formula (as opposed to a recursive formula) for the  $n$ th Fibonacci number by solving a difference equation.

- a) Let  $v_n = \begin{pmatrix} F_{n+1} \\ F_n \end{pmatrix}$ , so  $v_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ,  $v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ , etc. Find a state change matrix  $A$  such that  $v_{n+1} = Av_n$  for all  $n \geq 0$ .

- b) Show that the eigenvalues of  $A$  are  $\lambda_1 = \frac{1}{2}(1 + \sqrt{5})$  and  $\lambda_2 = \frac{1}{2}(1 - \sqrt{5})$ , with corresponding eigenvectors  $w_1 = \begin{pmatrix} -1 \\ \lambda_2 \end{pmatrix}$  and  $w_2 = \begin{pmatrix} -1 \\ \lambda_1 \end{pmatrix}$ .

[Hint: Check that  $Aw_i = \lambda_i w_i$  using the relations  $\lambda_1 \lambda_2 = -1$  and  $\lambda_1 + \lambda_2 = 1$ .]

- c) Expand  $v_0$  in this eigenbasis: that is, find  $x_1, x_2$  such that  $v_0 = x_1 w_1 + x_2 w_2$ . (It helps to write  $x_1, x_2$  in terms of  $\lambda_1, \lambda_2$ .)

- d) Multiply  $v_0 = x_1 w_1 + x_2 w_2$  by  $A^n$  to show that

$$F_n = \frac{\lambda_1^n - \lambda_2^n}{\lambda_1 - \lambda_2}.$$

- e) Use this formula to explain why  $F_{n+1}/F_n$  approaches the **golden ratio** when  $n$  is large.

5. Pretend that there are three **Red Box** kiosks in Durham. Let  $x_t, y_t, z_t$  be the number of copies of **Prognosis Negative** at each of the three kiosks, respectively, on day  $t$ . Suppose in addition that a customer renting a movie from kiosk  $i$  will return the movie the next day to kiosk  $j$ , with the following probabilities:

		Renting from kiosk		
		1	2	3
Returning to kiosk	1	30%	40%	50%
	2	30%	40%	30%
	3	40%	20%	20%

For instance, a customer renting from kiosk 3 has a 50% probability of returning it to kiosk 1.

- a) Let  $v_t = (x_t, y_t, z_t)$ . Find the state change matrix  $A$  such that  $v_{t+1} = Av_t$ .
- b) Diagonalize  $A$ . What are its eigenvalues?  
 [Hint:  $A$  is a stochastic matrix, so you know one eigenvalue by HW9#15(c).]
- c) If you start with a total of 1 000 copies of Prognosis Negative, how many of them will eventually end up at each kiosk? Does it matter what the initial state is?

This is an example of a **stochastic process**, and is an important application of eigenvalues and eigenvectors.

6. For each  $2 \times 2$  matrix  $A$  in Problem 1, if  $A$  is diagonalizable, find an invertible matrix  $C$  and a diagonal matrix  $D$  such that  $A = CDC^{-1}$ .
7. For each matrix  $A$  in Problem 2, if  $A$  is diagonalizable, find an invertible matrix  $C$  and a diagonal matrix  $D$  such that  $A = CDC^{-1}$ .
8. Consider the matrix

$$A = \begin{pmatrix} 4 & -3 & 0 \\ 2 & -1 & 0 \\ 1 & -1 & 1 \end{pmatrix}.$$

- a) Find a diagonal matrix  $D$  and an invertible matrix  $C$  such that  $A = CDC^{-1}$ .
- b) Find a *different* diagonal matrix  $D'$  and a *different* invertible matrix  $C'$  such that  $A = C'D'C'^{-1}$ .

[Hint: Try re-ordering the eigenvalues.]

9. Compute the matrix with eigenvalues 0, 1, 2 and corresponding eigenvectors

$$\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}.$$

(There is only one such matrix.)

10. Let  $A$  and  $B$  be  $n \times n$  matrices, and let  $v_1, \dots, v_n$  be a basis of  $\mathbf{R}^n$ .

a) Suppose that each  $v_i$  is an eigenvector of both  $A$  and  $B$ . Show that  $AB = BA$ .

b) Suppose that each  $v_i$  is an eigenvector of both  $A$  and  $B$  with the same eigenvalue. Show that  $A = B$ .

[Hint: Hint: use the matrix form of diagonalization.]

11. Let  $A$  be an  $n \times n$  matrix, and let  $C$  be an invertible  $n \times n$  matrix. Prove that the characteristic polynomial of  $CAC^{-1}$  equals the characteristic polynomial of  $A$ .

In particular,  $A$  and  $CAC^{-1}$  have the same eigenvalues, the same determinant, and the same trace. They are called *similar* matrices.

12. Let  $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ . Find a closed formula for  $A^n$ : that is, an expression of the form


$$A^n = \begin{pmatrix} a_{11}(n) & a_{12}(n) \\ a_{21}(n) & a_{22}(n) \end{pmatrix},$$

where  $a_{ij}(n)$  is a function of  $n$ .

13. A certain  $2 \times 2$  matrix  $A$  has eigenvalues 1 and 2. The eigenspaces are shown in the picture below.

a) Draw  $Av$ ,  $A^2v$ , and  $Aw$ .

b) Compute the limit of  $A^n v / \|A^n v\|$  as  $n \rightarrow \infty$ .



14. A certain diagonalizable  $2 \times 2$  matrix  $A$  is equal to  $CDC^{-1}$ , where  $C$  has columns  $w_1, w_2$  pictured below, and  $D = \begin{pmatrix} 1/3 & 0 \\ 0 & 1/2 \end{pmatrix}$ .

- Draw  $C^{-1}v$  on the left.
- Draw  $DC^{-1}v$  on the left.
- Draw  $Av = CDC^{-1}v$  on the right.
- What happens to  $A^n v$  as  $n \rightarrow \infty$ ?

