

KNOT AND BRAID INVARIANTS FROM CONTACT HOMOLOGY I

LENHARD NG

ABSTRACT. We introduce topological invariants of knots and braid conjugacy classes, in the form of differential graded algebras, and present an explicit combinatorial formulation for these invariants. The algebras conjecturally give the relative contact homology of certain Legendrian tori in five-dimensional contact manifolds. We present several computations and derive a relation between the knot invariant and the determinant.

1. INTRODUCTION

The study of invariants of Legendrian submanifolds in contact manifolds is currently a very active field of research. Holomorphic curve techniques stemming from the introduction of Symplectic Field Theory [7] have inspired a great deal of work on Legendrian isotopy invariants; papers in the subject include [3, 5, 6, 8, 9, 18, 23], among others. Most of these papers focus on Legendrian knots, and all deduce results in symplectic and contact geometry. Legendrian knots, however, form a restrictive class within the space of all knots, and Legendrian isotopy only makes sense in the presence of an ambient contact structure.

In this manuscript, we use ideas from Symplectic Field Theory to obtain new invariants of topological knots and braids. We note that holomorphic curves have been previously used to define knot invariants, through the work of Ozsváth and Szabó [21] on Heegaard Floer homology. Their approach, however, is very different from the one leading to the invariants in our paper, which is due to Eliashberg and can be briefly summarized as follows. A knot or braid yields in a natural way a Legendrian torus in a certain 5-dimensional contact manifold, and topological isotopy of the knot or braid results in Legendrian isotopy of the torus; thus, Legendrian invariants of the torus, and in particular relative contact homology, yield topological invariants for the original knot or braid.

We will discuss this correspondence further in Section 3, but the analytical underpinnings of the precise calculation of relative contact homology are still in progress. Instead, in the rest of the paper, we handle our new invariants algebraically, with no reference to contact geometry. The proofs of invariance, and all subsequent calculations, are completely combinatorial and algebraic.

We now describe our invariants. The starting point is a representation ϕ of the braid group B_n as a group of algebra automorphisms of a tensor algebra with $n(n-1)$ generators. This representation, which extends ones considered previously by Magnus [16] and Humphries [12], is nearly faithful, and there is a slightly larger representation ϕ^{ext} which is faithful. The map ϕ then gives rise to a differential graded algebra (customarily abbreviated “DGA” in the subject) for any braid B , which we call the braid DGA, while ϕ^{ext} yields a larger algebra called the knot DGA.

There is an equivalence relation on DGAs first introduced by Chekanov [3] called stable tame isomorphism, which has been the subject of much work because of its importance in the theory of Legendrian invariants. The main results of this paper state that, up to stable tame isomorphism, the braid DGA of B is an invariant of the conjugacy class of B , while the knot DGA of B is an invariant of the knot which is the closure of B .

In particular, the graded homology of these DGAs, which is unchanged by stable tame isomorphism, gives an invariant of braid conjugacy classes and knots; we call this the contact homology of the braid or knot. Although the homology of a DGA is generally hard to compute, there is a relatively simple form for the 0-dimensional part HC_0 of the contact homology, and we can compute HC_0 explicitly for a large number of knots.

We concentrate on the invariant HC_0 because it is difficult in general to distinguish between stable tame isomorphism classes of DGAs, and because HC_0 seems to encode much of the information from the knot or braid DGA. It is sometimes also hard to tell when two knots or braids have isomorphic HC_0 , but we can deduce numerical invariants from HC_0 , called augmentation numbers, which are easy to calculate by computer.

Knot contact homology seems to be an invariant distinct from previously known knot invariants. On the one hand, it does not distinguish between mirrors, and a result in [19] states that, for two-bridge knots, HC_0 depends only on the determinant $|\Delta_K(-1)|$ of the knot; hence one might expect that knot contact homology is fairly weak as an invariant. On the other hand, there are pairs of knots which are distinguished by HC_0 (via augmentation numbers) but not by (any given one of) a litany of familiar knot invariants: Alexander polynomial, Jones polynomial, HOMFLY polynomial, Kauffman polynomial, signature, Khovanov invariant, and Ozsváth–Szabó invariant.

One classical invariant that is contained in knot contact homology is the determinant; in fact, the determinant is encoded by a linearization HC_*^{lin} of HC_* . A further link between $HC_0(K)$ and the determinant is the existence of a surjection from $HC_0(K)$ to a ring $\mathbb{Z}[x]/(p(x))$, where p is a certain polynomial determined by the homology of the double branched cover over K ; this map is often an isomorphism.

Although knot contact homology is uniquely defined as a theory over \mathbb{Z}_2 , it has different viable lifts to \mathbb{Z} , one of which is the theory described above. However, there is another, inequivalent, lift to \mathbb{Z} , which we call alternate knot contact homology; the alternate theory lacks some of the properties of

the usual theory but produces a knot invariant which seems to contain more information.

The paper is organized as follows. In Section 2, we define the braid and knot DGAs and state the main invariance results. Section 3 gives a highly informal description of how our invariants relate to contact geometry. Section 4 defines contact homology for braids and knots and proves invariance of HC_0 ; we subsequently return to the invariance proof of the full DGAs in Section 5. Section 6 presents properties of the invariants, including augmentation numbers and behavior under mirrors, and Section 7 relates knot contact homology to the knot's determinant. Section 8 gives some examples of computations of the invariants. We discuss the alternate knot invariant in Section 9.

In the sequel to this paper [19], we give a geometric interpretation of ϕ via the usual treatment of B_n as a mapping class group. We then use this to present a new, completely topological definition for HC_0 for braids and knots, in terms of cords and relations similar to skein relations, and find a relation to character varieties. This interpretation explains, independent of contact geometry, “why” HC_0 is a topological invariant.

Acknowledgements. I am grateful to Yasha Eliashberg for suggesting the project and the means to approach it and offering a great deal of insightful guidance. I would also like to thank John Etnyre, Michael Hutchings, Kiran Kedlaya, Tom Mrowka, Josh Sabloff, Ravi Vakil, and Ke Zhu for useful conversations, and the Institute for Advanced Study, the American Institute of Mathematics, and Stanford University for their hospitality. Funding was provided by a Five-Year Fellowship from the American Institute of Mathematics, and at one point by NSF grant DMS-9729992.

2. DEFINITIONS AND MAIN RESULTS

2.1. Differential graded algebras. In contact geometry, the homology of a certain DGA defines the relative contact homology of a Legendrian submanifold in a contact manifold. In all known cases, a Legendrian isotopy invariant is provided not only by relative contact homology, but also by the stable tame isomorphism class of the DGA.

We recall the basic definitions, originally due to Chekanov [3]. Let $\mathbb{Z}\langle a_1, \dots, a_n \rangle$ denote the (noncommutative, unital) tensor algebra generated by a_1, \dots, a_n , which is generated as a \mathbb{Z} -module by words in the a_i 's, including the empty word.

Definition 2.1. Equip $\mathcal{A} = \mathbb{Z}\langle a_1, \dots, a_n \rangle$ with a grading over \mathbb{Z} by assigning degrees to the generators a_i . Then a *differential* on $\mathcal{A} = \mathbb{Z}\langle a_1, \dots, a_n \rangle$ is a map $\partial : \mathcal{A} \rightarrow \mathcal{A}$ which lowers degree by 1 and satisfies $\partial^2 = 0$ and the Leibniz rule $\partial(vw) = (\partial v)w + (-1)^{\deg v}v(\partial w)$. The pair (\mathcal{A}, ∂) is what we call a *differential graded algebra*, for the purposes of this paper.

To define a particular equivalence relation between DGAs known as stable tame isomorphism, we need several more notions, beginning with a refinement of the usual notion of isomorphism.

Definition 2.2. An algebra map $\phi: (\mathbb{Z}\langle a_1, \dots, a_n \rangle, \partial) \rightarrow (\mathbb{Z}\langle \tilde{a}_1, \dots, \tilde{a}_n \rangle, \tilde{\partial})$ is an *elementary isomorphism* if it is a graded chain map and, for some i ,

$$\phi(a_j) = \begin{cases} \pm \tilde{a}_i + v & \text{for } j = i \text{ and some } v \in \mathbb{Z}\langle \tilde{a}_1, \dots, \tilde{a}_{i-1}, \tilde{a}_{i+1}, \dots, \tilde{a}_n \rangle \\ \tilde{a}_j & \text{for } j \neq i. \end{cases}$$

A *tame isomorphism* between DGAs is any composition of elementary isomorphisms.

In the above definition, the order of the generators is immaterial; that is, a map which simply permutes the generators of the DGA (and treats the grading and differential accordingly) is considered to be elementary. Note also that elementary and tame isomorphisms are in fact isomorphisms.

A “trivial” example of a DGA is given by $(\mathcal{E}^i = \mathbb{Z}\langle e_1^i, e_2^i \rangle, \partial)$, where $\deg e_1^i - 1 = \deg e_2^i = i$ and $\partial e_1^i = e_2^i, \partial e_2^i = 0$. The main equivalence relation between DGAs, which we now present, stipulates that adding this trivial DGA does not change equivalence class.

Definition 2.3. The *degree- i algebraic stabilization* of the DGA $(\mathcal{A} = \mathbb{Z}\langle a_1, \dots, a_n \rangle, \partial)$ is the DGA $(S^i(\mathcal{A}) = \mathbb{Z}\langle a_1, \dots, a_n, e_1^i, e_2^i \rangle, \partial)$, with grading and differential induced from \mathcal{A} and \mathcal{E}^i . Two DGAs (\mathcal{A}, ∂) and $(\tilde{\mathcal{A}}, \tilde{\partial})$ are *stable tame isomorphic* if there exist algebraic stabilizations $(S^{i_1}(\dots(S^{i_k}(\mathcal{A}))\dots), \partial)$ and $(S^{\tilde{i}_1}(\dots(S^{\tilde{i}_k}(\tilde{\mathcal{A}}))\dots), \tilde{\partial})$ which are tamely isomorphic.

Stable tame isomorphism is designed to be a special case of quasi-isomorphism.

Proposition 2.4. *If (\mathcal{A}, ∂) and $(\tilde{\mathcal{A}}, \tilde{\partial})$ are stable tame isomorphic, then the homologies $H_*(\mathcal{A}, \partial)$ and $H_*(\tilde{\mathcal{A}}, \tilde{\partial})$ are isomorphic.*

The proof of Proposition 2.4 can be found, e.g., in [9].

2.2. The invariants. We now introduce the knot and braid invariants. The definition of the knot DGA depends on the braid definition, which we give first.

Let B_n denote the braid group on n strands, which is generated by $\sigma_1, \dots, \sigma_{n-1}$, with relations $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ for $1 \leq i \leq n-1$, and $\sigma_i \sigma_j = \sigma_j \sigma_i$ for $|i-j| \geq 2$. Write \mathcal{A}_n for the tensor algebra on $n(n-1)$ generators which we label as a_{ij} for $1 \leq i, j \leq n$ and $i \neq j$, and let $\text{Aut}(\mathcal{A}_n)$ denote the group of algebra automorphisms of \mathcal{A}_n .

For each generator σ_k of B_n , define the tame automorphism $\phi_{\sigma_k} \in \text{Aut}(\mathcal{A}_n)$ by its action on the generators of \mathcal{A}_n :

$$\phi_{\sigma_k} : \begin{cases} a_{ki} & \mapsto -a_{k+1,i} - a_{k+1,k}a_{ki} & i \neq k, k+1 \\ a_{ik} & \mapsto -a_{i,k+1} - a_{ik}a_{k,k+1} & i \neq k, k+1 \\ a_{k+1,i} & \mapsto a_{ki} & i \neq k, k+1 \\ a_{i,k+1} & \mapsto a_{ik} & i \neq k, k+1 \\ a_{k,k+1} & \mapsto a_{k+1,k} \\ a_{k+1,k} & \mapsto a_{k,k+1} \\ a_{ij} & \mapsto a_{ij} & i, j \neq k, k+1. \end{cases}$$

Proposition 2.5. ϕ extends to a homomorphism from B_n to $\text{Aut}(\mathcal{A}_n)$.

Proof. The fact that ϕ preserves the braid relations in B_n is a direct computation. \square

We will denote the image of $B \in B_n$ under ϕ as $\phi_B \in \text{Aut}(\mathcal{A}_n)$.

This representation was essentially first studied by Magnus [16] in the context of automorphisms of free groups; more precisely, he presented a representation on a polynomial algebra in $\binom{n}{2}$ variables, which we can derive from ϕ by abelianizing and setting $a_{ij} = a_{ji}$ for all i, j . Humphries [12] subsequently extended the representation to a polynomial algebra in $n(n-1)$ variables, and interpreted it in terms of transvections. Our homomorphism is simply the lift of Humphries' representation to the corresponding tensor algebra. We note that the starting point in [19] is a new topological interpretation of ϕ in terms of skein relations.

We can obtain a linear representation from ϕ by finding a finite-dimensional piece of \mathcal{A}_n on which ϕ descends; in the language of [8], this involves linearizing ϕ with respect to some augmentation. Concretely, if we write \mathcal{M}_n as the subalgebra of \mathcal{A}_n generated by $\{a_{ij} + 2\}$, then ϕ descends to $\mathcal{M}_n/\mathcal{M}_n^2$. If we further quotient by setting $a_{ij} = a_{ji}$ for all i, j , then we obtain an $\binom{n}{2}$ -dimensional representation of B_n , which is precisely the inverse of the Lawrence–Krammer representation [14] with $q = -1$ and $t = 1$. Indeed, there is a striking similarity in appearance between ϕ and the Lawrence–Krammer representation which we further explore in [19].

Unlike Lawrence–Krammer, ϕ does not give a faithful representation of B_n ; this is not overly surprising, since B_n has nontrivial center, while $\text{Aut}(\mathcal{A}_n)$ does not. Indeed, a result from [12] states that the kernel of ϕ is precisely the center of B_n , which is generated by $(\sigma_1 \cdots \sigma_{n-1})^n$. However, the extension map $\phi^{\text{ext}}: B_n \hookrightarrow B_{n+1} \xrightarrow{\phi} \text{Aut}(\mathcal{A}_{n+1})$ is faithful, a fact first established algebraically in [16]. We will return to ϕ^{ext} in a moment; geometric proofs of the faithfulness results can be found in [19].

We now define the DGA invariant for braids.

Definition 2.6. Let $B \in B_n$ be a braid. Let \mathcal{A} be the graded tensor algebra on $2n(n-1)$ generators, $\{a_{ij} \mid 1 \leq i, j \leq n, i \neq j\}$ of degree 0 and

$\{b_{ij} \mid 1 \leq i, j \leq n, i \neq j\}$ of degree 1. Define the differential ∂ on \mathcal{A} by

$$\partial b_{ij} = a_{ij} - \phi_B(a_{ij}), \quad \partial a_{ij} = 0.$$

Then we call (\mathcal{A}, ∂) the *braid DGA* of B .

Theorem 2.7. *If B, \tilde{B} are conjugate in B_n , then their braid DGAs are stable tame isomorphic.*

We defer the proof of Theorem 2.7 until Section 5.1. The DGA invariant for knots is also derived from the map ϕ ; in fact, we will see in Section 4.2 that it contains the braid DGA. Before introducing it, we need some more notation.

As mentioned before, we can define a map $\phi^{\text{ext}}: B_n \rightarrow \text{Aut}(\mathcal{A}_{n+1})$ induced from ϕ through the inclusion $B_n \hookrightarrow B_{n+1}$ obtained by adding an $(n+1)$ -st strand which does not interact with the other n strands; that is, each generator $\sigma_i \in B_n$ is mapped to $\sigma_i \in B_{n+1}$ for $1 \leq i \leq n-1$. When dealing with ϕ^{ext} , we replace all indices $(n+1)$ by an asterisk; i.e., write a_{*i}, a_{i*} for $a_{n+1,i}, a_{i,n+1}$, respectively. Since the $(n+1)$ -st strand does not interact with the other strands, it follows from the definition of ϕ that for $B \in B_n$, $\phi_B^{\text{ext}}(a_{*i})$ is a linear combination of a_{*j} , $1 \leq j \leq n$, with coefficients in \mathcal{A}_n , and similarly for $\phi_B^{\text{ext}}(a_{i*})$.

Definition 2.8. The $n \times n$ matrices $\Phi_B^L(A)$ and $\Phi_B^R(A)$ are defined by

$$\phi_B^{\text{ext}}(a_{i*}) = \sum_{j=1}^n (\Phi_B^L(A))_{ij} a_{j*} \quad \text{and} \quad \phi_B^{\text{ext}}(a_{*j}) = \sum_{i=1}^n a_{*i} (\Phi_B^R(A))_{ij}.$$

The matrices $\Phi_B^L(A)$ and $\Phi_B^R(A)$ can be obtained from each other by the “conjugation” operation discussed in Section 6.1. Though $\Phi_B^L(A), \Phi_B^R(A)$ are derived from ϕ , they also determine ϕ ; see Proposition 6.2.

To define the DGA for knots, it is convenient to assemble generators of the DGA into matrices. For notational purposes, let A denote the matrix (a_{ij}) , where we set $a_{ii} = -2$ for all i , and similarly write $B = (b_{ij}), C = (c_{ij}), D = (d_{ij})$, where b_{ij}, c_{ij}, d_{ij} are variables used below with no stipulation on b_{ii}, c_{ii}, d_{ii} ; also, write ∂A for the matrix (∂a_{ij}) , $\phi_B(A)$ for $(\phi_B(a_{ij}))$, and so forth. (There should hopefully be no confusion between a braid B and the matrix B .) We will sometimes view A as a matrix of variables, so that, e.g., we define $\Phi_B^L(M)$ for an $n \times n$ matrix M to be the evaluation of $\Phi_B^L(A)$ when we set $a_{ij} = M_{ij}$ for all i, j . (Note that this requires $M_{ii} = -2$ for all i .)

Definition 2.9. Let $B \in B_n$. Write \mathcal{A} for the tensor algebra with the following generators: $\{a_{ij} \mid 1 \leq i, j \leq n, i \neq j\}$ of degree 0; $\{b_{ij}, c_{ij} \mid 1 \leq i, j \leq n\}$ of degree 1; and $\{d_{ij} \mid 1 \leq i, j \leq n\}$ and $\{e_i \mid 1 \leq i \leq n\}$ of degree

2. Define the differential ∂ on \mathcal{A} by

$$\begin{aligned}\partial A &= 0 \\ \partial B &= (1 - \Phi_B^L(A)) \cdot A \\ \partial C &= A \cdot (1 - \Phi_B^R(A)) \\ \partial D &= B \cdot (1 - \Phi_B^R(A)) - (1 - \Phi_B^L(A)) \cdot C \\ \partial e_i &= (B + \Phi_B^L(A) \cdot C)_{ii},\end{aligned}$$

where \cdot denotes matrix multiplication. Then we call (\mathcal{A}, ∂) the *knot DGA* of B .

The fact that $\partial^2 = 0$ in the knot DGA is not obvious; while it is clear that $\partial^2 A = \partial^2 B = \partial^2 C = \partial^2 D = 0$, we have $\partial^2 e_i = -2 - (\Phi_B^L(A) \cdot A \cdot \Phi_B^R(A))_{ii}$, and we need Proposition 4.7 to conclude that $\partial^2 e_i = 0$. The key result for knot DGAs is the following theorem, which allows the knot DGA to descend from braids to knots.

Theorem 2.10. *If the closures of two braids B, \tilde{B} are the same knot, then the knot DGAs for B, \tilde{B} are stable tame isomorphic.*

The proof of Theorem 2.10 is postponed until Section 5.2, after we first introduce HC_0 and present the easier proof of invariance for HC_0 .

By Alexander's Theorem, every knot in \mathbb{R}^3 can be represented as a closed braid. Theorem 2.10 implies that the following notion is well-defined.

Definition 2.11. Let $K \subset \mathbb{R}^3$ be a knot. The *knot DGA class* of K is the equivalence class (under stable tame isomorphism) of the knot DGA of any braid whose closure is K .

We will occasionally abuse notation and speak of the ‘‘knot DGA’’ of K ; this is understood to be modulo equivalence. The knot DGA can equally well be defined for links, for which it also gives an invariant.

We conclude this section by explicitly computing, as an example, the knot DGA for the trefoil, which is the closure of $\sigma_1^3 \in B_2$. By the definition of ϕ , we see that $\phi_{\sigma_1}^{\text{ext}}$ acts as follows: $a_{12} \mapsto a_{21}$, $a_{21} \mapsto a_{12}$, $a_{1*} \mapsto -a_{2*} - a_{21}a_{1*}$, $a_{2*} \mapsto a_{1*}$, $a_{*1} \mapsto -a_{*2} - a_{*1}a_{12}$, $a_{*2} \mapsto a_{*1}$. Iterating $\phi_{\sigma_1}^{\text{ext}}$ three times on $a_{1*}, a_{2*}, a_{*1}, a_{*2}$ yields

$$\Phi_{\sigma_1^3}^L(A) = \begin{pmatrix} 2a_{21} - a_{21}a_{12}a_{21} & 1 - a_{21}a_{12} \\ -1 + a_{12}a_{21} & a_{12} \end{pmatrix} \quad \text{and} \quad \Phi_{\sigma_1^3}^R(A) = \begin{pmatrix} 2a_{12} - a_{12}a_{21}a_{12} & -1 + a_{12}a_{21} \\ 1 - a_{21}a_{12} & a_{21} \end{pmatrix}.$$

One can now calculate the differential on the knot DGA of σ_1^3 :

$$\begin{aligned}
\partial b_{11} &= -2 + 3a_{21} - a_{21}a_{12}a_{21} & \partial c_{11} &= -2 + 3a_{12} - a_{12}a_{21}a_{12} \\
\partial b_{12} &= 2 + a_{12} - 4a_{21}a_{12} + a_{21}a_{12}a_{21}a_{12} & \partial c_{21} &= 2 + a_{21} - 4a_{21}a_{12} + a_{21}a_{12}a_{21}a_{12} \\
\partial b_{21} &= -2 + a_{21} + a_{12}a_{21} & \partial c_{12} &= -2 + a_{12} + a_{12}a_{21} \\
\partial b_{22} &= -2 + 3a_{12} - a_{12}a_{21}a_{12} & \partial c_{22} &= -2 + 3a_{21} - a_{21}a_{12}a_{21} \\
\partial d_{11} &= b_{11} - b_{12} - c_{11} + c_{21} + 2a_{21}c_{11} - 2b_{11}a_{12} - a_{21}a_{12}c_{21} + b_{12}a_{21}a_{12} \\
&\quad - a_{21}a_{12}a_{21}c_{11} + b_{11}a_{12}a_{21}a_{12} \\
\partial d_{12} &= b_{11} + b_{12} - c_{12} + c_{22} + 2a_{21}c_{12} - b_{12}a_{21} - a_{21}a_{12}c_{22} \\
&\quad - b_{11}a_{12}a_{21} - a_{21}a_{12}a_{21}c_{12} \\
\partial d_{21} &= b_{21} - b_{22} - c_{11} - c_{21} + a_{12}c_{21} - 2b_{21}a_{12} + a_{12}a_{21}c_{11} \\
&\quad + b_{22}a_{21}a_{12} + b_{21}a_{12}a_{21}a_{12} \\
\partial d_{22} &= b_{21} + b_{22} - c_{12} - c_{22} + a_{12}c_{22} - b_{22}a_{21} + a_{12}a_{21}c_{12} - b_{21}a_{12}a_{21} \\
\partial e_1 &= b_{11} + c_{21} + 2a_{21}c_{11} - a_{21}a_{12}c_{21} - a_{21}a_{12}a_{21}c_{11} \\
\partial e_2 &= b_{22} - c_{12} + a_{12}c_{22} + a_{12}a_{21}c_{12}.
\end{aligned}$$

We will return to this example in future sections.

3. MOTIVATION FROM CONTACT GEOMETRY

Having defined the knot and braid DGAs in the previous section, we now describe the background in contact geometry leading to the development of these invariants. It should be noted that work is in progress to establish rigorously that the DGAs give the relative contact homology theory which we describe below.

Given a smooth manifold M , the cotangent bundle T^*M has a canonical symplectic structure given by $\omega = d\lambda$, where λ is the tautological one-form on T^*M sending a tangent vector in T^*M to the pairing between the base point in T^*M and the projected tangent vector in TM under the projection $\pi : T^*M \rightarrow M$. For any smooth submanifold $N \subset M$, the conormal bundle $\mathcal{L}N = \{\theta \in T^*M \mid \langle \theta, v \rangle = 0 \text{ for all } v \in T_{\pi(v)}N\}$ over N is then a Lagrangian submanifold of T^*M , and smooth isotopy of N leads to Lagrangian isotopy of $\mathcal{L}N$. This setup (with N a knot) was introduced in the physics literature in [20], and was communicated to the author in this generality by Eliashberg.

Partly because $\mathcal{L}N$ is noncompact, it is more convenient for us to consider a slightly modified setup. Suppose that M is equipped with a metric. Then the cosphere bundle ST^*M of unit covectors in T^*M (“co-oriented contact elements in M ”) has a canonical contact structure induced by the one-form λ . If $N \subset M$ is a compact smooth submanifold, then the unit conormal bundle $LN = \mathcal{L}N \cap ST^*M$ is a compact Legendrian submanifold of ST^*M , and smooth isotopy of N leads to Legendrian isotopy of LN .

In recent years, beginning with work of Eliashberg and Hofer, invariants of Legendrian isotopy in contact manifolds have been developed, via holomorphic-curve techniques, using relative contact homology [6] and the

more general Symplectic Field Theory [7]. In particular, for Legendrian knots in \mathbb{R}^3 with the standard contact structure, Chekanov [3] developed a purely combinatorial formulation of relative contact homology; this work was subsequently extended in [9, 18]. Relative contact homology was also studied for knots in circle bundles in [23], and for \mathbb{R}^{2n+1} with the standard contact structure in [5]. In all of these special cases, relative contact homology is given by the homology of a certain DGA whose stable tame isomorphism class is an invariant. With the exception of [5], however, the DGAs are defined combinatorially rather than geometrically, with invariance proofs given by combinatorics as well. The invariants in the present paper are likewise combinatorial in nature.

We now specialize, as in [20], to the case when $M = \mathbb{R}^3$ and $K \subset \mathbb{R}^3$ is a knot. Then LK is an embedded Legendrian 2-torus in the 5-dimensional contact manifold $ST^*\mathbb{R}^3 = \mathbb{R}^3 \times S^2$, which is contactomorphic to the 1-jet space $J^1(S^2)$ with the standard contact structure. We note that LK is always topologically unknotted since it has codimension 3 in $ST^*\mathbb{R}^3$; however, its Legendrian isotopy type yields information about the knot K . In fact, the knot DGA defined in Section 2.2, which is a nontrivial invariant, conjecturally gives the relative contact homology of LK in $ST^*\mathbb{R}^3$.

A local model suggested by Eliashberg allows us to study conjugacy classes of braids as a first step. Let $K \subset \mathbb{R}^3$ be any knot (or, in particular, the unknot), and let B be a braid. A sufficiently small neighborhood of LK is contactomorphic to $J^1(LK) = J^1(T^2)$. If we glue B along a tubular neighborhood of K to produce a new knot \tilde{K} , then $L\tilde{K}$ is in an arbitrarily small neighborhood of LK , and thus gives a Legendrian 2-torus in $J^1(T^2)$. A smooth isotopy of B , or a conjugation operation on B , yields a Legendrian isotopy of this torus in $J^1(T^2)$. Then the braid DGA from Section 2.2 conjecturally gives the relative contact homology of $L\tilde{K}$ in $J^1(T^2)$.

Recall from, e.g., [6] that the relative contact homology for a Legendrian submanifold L of a contact manifold (V, α) is the homology of a DGA whose generators are *Reeb chords* of L , i.e., flow lines of the Reeb vector field which begin and end on L . The differential is then defined by counting certain holomorphic maps of punctured disks into the symplectization $V \times \mathbb{R}$ of V , with boundary on $L \times \mathbb{R}$ and punctures limiting to Reeb chords.

For the case of a knot K in \mathbb{R}^3 , the contact manifold is $\mathbb{R}^3 \times S^2$, with Reeb vector field pointing in the \mathbb{R}^3 fibers, in the direction specified by the underlying point in S^2 . It follows that Reeb chords for LK correspond to “binormal chords” to K in \mathbb{R}^3 , i.e., oriented line segments in \mathbb{R}^3 with endpoints on K which are normal to K at both endpoints. For instance, the unknot given by an ellipse in the plane yields four Reeb chords, corresponding to the major and minor axes traversed either way.

If K is the closure of a braid $B \in B_n$, we can embed B in a neighborhood of an elliptical unknot in such a way that the Reeb chords for LK come in two families:

- $2n(n-1)$ “small chords” within the neighborhood of the unknot, four for each pair of braid strands; these divide further into two Reeb chords where the pair of strands is closest (a_{ij}) and two where the pair is farthest (b_{ij})
- $4n^2$ “big chords” corresponding to the Reeb chords of the ellipse; each binormal chord of the ellipse yields n^2 binormal chords for K (c_{ij}, d_{ij} for the minor-axis chords, e_{ij}, f_{ij} for the major-axis chords), since there are n choices for each endpoint.

For the braid itself, the Reeb chords for $LB \subset J^1(T^2)$ correspond to the $2n(n-1)$ “small chords” a_{ij}, b_{ij} .

This gives a rough explanation for the generators of the knot and braid DGAs; for the knot case, this actually gives the modified knot DGA defined below in Proposition 4.8. Showing that the differentials of these DGAs actually count the appropriate holomorphic disks is beyond the scope of this paper. The analytical details, which use an approach due to Fukaya and Oh [10] of counting gradient flow trees, are the subject of work in progress. As in the theory of Legendrian knots in standard contact \mathbb{R}^3 , however, the combinatorial proof of the invariance of our DGAs under isotopy gives evidence for the validity of our “computation” of relative contact homology.

4. KNOT AND BRAID CONTACT HOMOLOGY

In this section, we examine the homology of the DGAs defined in Section 2, and focus on a particular piece for which isotopy invariance is easy to establish. We will return to invariance proofs of the full DGAs in Section 5.

4.1. Definitions. In order to use the knot and braid DGAs as invariants, we need to know when two DGAs are stable tame isomorphic. This can be done through “computable” invariants of equivalence classes of DGAs, including Poincaré polynomials [3] and the characteristic algebra [18].

In the case of braid and knot DGAs, it turns out that the first order Poincaré polynomial over \mathbb{Z}_2 corresponding to the trivial augmentation gives no interesting invariant of the braid conjugacy class or the knot. (See however Section 8.3 for a different result for a nontrivial augmentation over \mathbb{Z}_2 , and Section 7.2 for linearization over \mathbb{Z} .) This is because to first order over \mathbb{Z}_2 , the homomorphism ϕ factors through the projection $B_n \rightarrow S_n$ to the symmetric group. However, the degree 0 part of the characteristic algebra already yields interesting invariants; we recall from [18] that the characteristic algebra of a DGA (\mathcal{A}, ∂) is the quotient of \mathcal{A} by the two-sided ideal generated by $\text{im } \partial$. Before proceeding further, we define the contact homology of a braid or knot.

Definition 4.1. The *contact homology* of a braid B , written $HC_*(B)$, is defined to be the (graded) homology of the braid DGA of B . The *contact homology* of a knot K , written $HC_*(K)$, is the (graded) homology of the

knot DGA of any braid whose closure is K . More generally, if R is any ring, then we define as usual $HC_*(B; R)$ and $HC_*(K; R)$ to be the homology of the appropriate DGA tensored with R .

Proposition 4.2. *The degree 0 contact homology of knots and braids is the degree 0 part of the characteristic algebra of the corresponding DGA. More explicitly, for $B \in B_n$, let $\mathcal{I}_B^{\text{braid}}$ be the two-sided ideal of \mathcal{A}_n generated by $\{a_{ij} - \phi_B(a_{ij})\}$, while $\mathcal{I}_B^{\text{knot}}$ is the two-sided ideal of \mathcal{A}_n generated by the entries of the two matrices $(1 - \Phi_B^L(A)) \cdot A$ and $A \cdot (1 - \Phi_B^R(A))$. Then $HC_0(B) = \mathcal{A}_n / \mathcal{I}_B^{\text{braid}}$ and $HC_0(K) = \mathcal{A}_n / \mathcal{I}_B^{\text{knot}}$, where K is the closure of B .*

Proof. Since the braid and knot DGAs of $B \in B_n$ have no generators in negative degree, the entirety of \mathcal{A}_n (which is the degree 0 part of the algebra) consists of cycles. On the other hand, the image of ∂ in degree 0 is precisely the ideal generated by the images under ∂ of the generators in degree 1, and so the ideal of boundaries in degree 0 is given by $\mathcal{I}_B^{\text{braid}}$, $\mathcal{I}_B^{\text{knot}}$ for the braid and knot DGA, respectively. \square

For example, for the trefoil, $HC_0(3_1)$ is given by $\mathbb{Z}\langle a_{12}, a_{21} \rangle$ modulo the relations provided by $\partial b_{ij}, \partial c_{ij}$ as calculated in Section 2.2. From ∂b_{21} and ∂c_{12} , we have relations $-2 + a_{21} + a_{12}a_{21} = -2 + a_{12} + a_{12}a_{21} = 0$, and hence $a_{12} = a_{21}$. If we set $a_{12} = a_{21} = x$, then the $\partial b_{ij}, \partial c_{ij}$ relations become $\{x^3 - 3x + 2, x^4 - 4x^2 + x + 2, x^2 + x - 2\}$. The gcd of these polynomials is $x^2 + x - 2$, and so $HC_0(3_1) \cong \mathbb{Z}[x]/(x^2 + x - 2)$.

In practice, we will study HC_0 rather than the full braid and knot DGAs, because it is relatively easy to compute, as the trefoil example demonstrates. It is immediate from Proposition 2.4 and Theorems 2.6 and 2.9 that HC_0 gives an invariant of braid group conjugacy classes and knots. However, there is a direct proof of the invariance of HC_0 which is much simpler notationally than the proofs of Theorems 2.6 and 2.9, while containing the main ideas from these proofs. We will give this direct proof below for braids, and in Section 4.3 for knots.

Theorem 4.3. *Up to isomorphism, $HC_0(B)$ is an invariant of the conjugacy class of the braid B .*

Proof. Suppose that B, \tilde{B} are conjugate in the braid group B_n , so that $\tilde{B} = C^{-1}BC$ for some $C \in B_n$. We use the notation of Proposition 4.2.

First note that $\mathcal{I}_B^{\text{braid}} \subset \mathcal{A}_n$ is the ideal generated by the image of the map $1 - \phi_B: \mathcal{A}_n \rightarrow \mathcal{A}_n$: if $x_1, x_2 \in \mathcal{A}_n$, then $(1 - \phi_B)(x_1x_2) = (1 - \phi_B)(x_1)x_2 + \phi_B(x_1)(1 - \phi_B)(x_2)$ is in the ideal generated by $(1 - \phi_B)(x_1)$ and $(1 - \phi_B)(x_2)$. Similarly, $\mathcal{I}_{\tilde{B}}^{\text{braid}}$ is the ideal generated by the image of $1 - \phi_{\tilde{B}} = 1 - \phi_C^{-1}\phi_B\phi_C$.

It follows by inspection of the commutative diagram

$$\begin{array}{ccc} \mathcal{A}_n & \xrightarrow{1-\phi_B} & \mathcal{A}_n \\ \phi_C \downarrow \cong & & \cong \downarrow \phi_C \\ \mathcal{A}_n & \xrightarrow{1-\phi_C^{-1}\phi_B\phi_C} & \mathcal{A}_n \end{array}$$

that $\mathcal{I}_{\tilde{B}}^{\text{braid}}$ is the image of $\mathcal{I}_B^{\text{braid}}$ under the automorphism ϕ_C on \mathcal{A}_n , and hence that $HC_0(B) = \mathcal{A}_n/\mathcal{I}_B^{\text{braid}}$ and $HC_0(\tilde{B}) = \mathcal{A}_n/\mathcal{I}_{\tilde{B}}^{\text{braid}}$ are isomorphic. \square

The corresponding invariance result for knots requires a few preparatory results, which are interesting in their own right; we give these in the following section, and prove HC_0 invariance for knots in Section 4.3.

4.2. Ancillary results. Here we collect a number of results about ϕ , Φ^L , and Φ^R which will be crucial for future sections.

Proposition 4.4 (Chain Rule). *If $B_1, B_2 \in B_n$, then*

$$\Phi_{B_1 B_2}^L(A) = \Phi_{B_2}^L(\phi_{B_1}(A)) \cdot \Phi_{B_1}^L(A) \quad \text{and} \quad \Phi_{B_1 B_2}^R(A) = \Phi_{B_1}^R(A) \cdot \Phi_{B_2}^R(\phi_{B_1}(A)).$$

Proof. We will establish the result for Φ^L ; the proof for Φ^R is completely analogous. By the definition of Φ^L , we have

$$\phi_{B_1 B_2}^{\text{ext}}(a_{i*}) = \sum_{j=1}^n (\Phi_{B_1 B_2}^L(A))_{ij} a_{j*};$$

on the other hand,

$$\begin{aligned} \phi_{B_1 B_2}^{\text{ext}}(a_{i*}) &= \phi_{B_1}^{\text{ext}} \phi_{B_2}^{\text{ext}}(a_{i*}) \\ &= \sum_{j=1}^n \phi_{B_1}^{\text{ext}} \left((\Phi_{B_2}^L(A))_{ij} a_{j*} \right) \\ &= \sum_{j=1}^n (\Phi_{B_2}^L(\phi_{B_1}(A)))_{ij} \phi_{B_1}^{\text{ext}}(a_{j*}) \\ &= \sum_{j,\ell=1}^n (\Phi_{B_2}^L(\phi_{B_1}(A)))_{ij} (\Phi_{B_1}^L(A))_{j\ell} a_{\ell*}, \end{aligned}$$

and the result follows.

A word of explanation is in order for the penultimate equality in the chain above, because it may seem backwards. Applying $\phi_{B_1}^{\text{ext}}$ to $(\Phi_{B_2}^L(A))_{ij} a_{j*}$ entails replacing each a_{ij} by $\phi_{B_1}(a_{ij})$ and each a_{j*} by $\phi_{B_1}^{\text{ext}}(a_{j*})$, and the result is $(\Phi_{B_2}^L(\phi_{B_1}(A)))_{ij} \phi_{B_1}^{\text{ext}}(a_{j*})$. \square

Corollary 4.5. *For any braid B , $\Phi_{B^{-1}}^L(\phi_B(A)) = (\Phi_B^L(A))^{-1}$ and $\Phi_{B^{-1}}^R(\phi_B(A)) = (\Phi_B^R(A))^{-1}$.*

Proof. Set $B_1 = B$, $B_2 = B^{-1}$ in Proposition 4.4, and use the fact that $\Phi_{\text{id}}^L(A) = \Phi_{\text{id}}^R(A) = 1$. \square

Lemma 4.6. *The matrices $\Phi_{\sigma_k}^L(A)$ and $\Phi_{\sigma_k}^R(A)$ are identical to the $n \times n$ identity matrix, outside of the 2×2 submatrix formed by rows $k, k+1$ and columns $k, k+1$, which is given by $\begin{pmatrix} -a_{k+1,k} & -1 \\ 1 & 0 \end{pmatrix}$ for $\Phi_{\sigma_k}^L(A)$ and $\begin{pmatrix} -a_{k,k+1} & 1 \\ -1 & 0 \end{pmatrix}$ for $\Phi_{\sigma_k}^R(A)$.*

Proof. This is immediate from the definitions of ϕ , Φ^L , and Φ^R . \square

Proposition 4.7. *For any braid B , $\phi_B(A) = \Phi_B^L(A) \cdot A \cdot \Phi_B^R(A)$.*

Proof. When $B = \sigma_k$ for some k , the identity can be verified by direct calculation, using Lemma 4.6. We conclude that the identity also holds for $B = \sigma_k^{-1}$, either by another direct calculation, or by using the result for $B = \sigma_k$: since $\phi_{\sigma_k}(A) = \Phi_{\sigma_k}^L(A) \cdot A \cdot \Phi_{\sigma_k}^R(A)$, we have

$$A = \Phi_{\sigma_k}^L(\phi_{\sigma_k}^{-1}(A)) \cdot \phi_{\sigma_k}^{-1}(A) \cdot \Phi_{\sigma_k}^R(\phi_{\sigma_k}^{-1}(A)) = \left(\Phi_{\sigma_k}^L(A) \right)^{-1} \cdot \phi_{\sigma_k}^{-1}(A) \cdot \left(\Phi_{\sigma_k}^R(A) \right)^{-1},$$

where the last equality comes from Corollary 4.5; the desired identity for $B = \sigma_k^{-1}$ follows.

We now show that if the identity holds for $B = B_1$ and $B = B_2$, then it holds for $B = B_1 B_2$; the proposition then follows by induction. Indeed, assuming the identity for $B = B_1$ and $B = B_2$, we have

$$\begin{aligned} \phi_{B_1 B_2}(A) &= \phi_{B_1}(\phi_{B_2}(A)) \\ &= \Phi_{B_2}^L(\phi_{B_1}(A)) \cdot \phi_{B_1}(A) \cdot \Phi_{B_2}^R(\phi_{B_1}(A)) \\ &= \Phi_{B_2}^L(\phi_{B_1}(A)) \cdot \Phi_{B_1}^L(A) \cdot A \cdot \Phi_{B_1}^R(A) \cdot \Phi_{B_2}^R(\phi_{B_1}(A)) \\ &= \Phi_{B_1 B_2}^L(A) \cdot A \cdot \Phi_{B_1 B_2}^R(A), \end{aligned}$$

where we use the chain rule for the final equality. \square

A more geometric proof of Proposition 4.7 is given in [19]. As remarked in Section 2.2, Proposition 4.7 implies that the knot DGA does in fact satisfy $\partial^2 = 0$. It also follows from Proposition 4.7 that the full homomorphism ϕ_B can be deduced from the matrices Φ_B^L and Φ_B^R , which in turn can be deduced from each other via the conjugation map to be discussed in Section 6.1. Thus we may view Φ_B^L and Φ_B^R as “square roots” of ϕ_B .

One final result we will need in the future is a reformulation of the knot DGA in terms of an equivalent DGA which has more generators but a slightly simpler definition.

Proposition 4.8. *Let $B \in B_n$. The knot DGA for B is stable tame isomorphic to the “modified knot DGA” which has generators $\{a_{ij} \mid 1 \leq i, j \leq n, i \neq j\}$ of degree 0, $\{b_{ij} \mid 1 \leq i, j \leq n, i \neq j\}$ and $\{c_{ij}, d_{ij} \mid 1 \leq i, j \leq n\}$ of*

degree 1, and $\{e_{ij}, f_{ij} \mid 1 \leq i, j \leq n\}$ of degree 2, with differential

$$\begin{aligned}\partial A &= 0 \\ \partial B &= A - \phi_B(A) \\ \partial C &= (1 - \Phi_B^L(A)) \cdot A \\ \partial D &= A \cdot (1 - \Phi_B^R(A)) \\ \partial E &= B - D - C \cdot \Phi_B^R(A) \\ \partial F &= B - C - \Phi_B^L(A) \cdot D.\end{aligned}$$

Here we set $b_{ii} = 0$ for all i .

Proof. In the modified knot DGA, replace e_{ij} by $e_{ij} + f_{ij}$ for all i, j ; after applying the induced tame automorphism, we obtain the same differential, except with $\partial E = C \cdot (1 - \Phi_B^R(A)) - (1 - \Phi_B^L(A)) \cdot D$. Now replace b_{ij} by $(B + C + \Phi_B^L(A) \cdot D)_{ij}$ for $i \neq j$, which constitutes another tame automorphism; the differential is again unchanged, except that $\partial B = 0$ (this follows from Proposition 4.7) and $\partial F = B - \text{diag}(C + \Phi_B^L(A) \cdot D)$, where diag replaces all non-diagonal entries by 0. Now we have $\partial f_{ij} = b_{ij}$ for $i \neq j$, and hence, up to stabilization, we may drop the generators $\{b_{ij}, f_{ij} \mid i \neq j\}$ from the DGA. The resulting DGA is precisely the knot DGA for B , after we relabel $c_{ij}, d_{ij}, e_{ij}, f_{ii}$ by $b_{ij}, c_{ij}, d_{ij}, -e_i$, respectively. \square

An important consequence of Proposition 4.8 is that the knot DGA for a braid can be obtained from the braid DGA, up to stable tame isomorphism, by adding generators and appropriately extending the definition of the differential.

Corollary 4.9. *If K is the closure of B , then $HC_0(K)$ is a quotient of $HC_0(B)$.*

Proof. Immediate from Propositions 2.4 and 4.8. \square

4.3. Invariance of HC_0 for knots. In this section, we prove the following result.

Theorem 4.10. *Up to isomorphism, $HC_0(K)$ is a well-defined invariant of a knot K .*

Proof. If B is a braid, we temporarily write $HC_0^{\text{knot}}(B)$ for the 0-dimensional homology of the knot DGA of B , as expressed in Proposition 4.2. We wish to show that if B, \tilde{B} are two braids whose closure is the same knot, then $HC_0^{\text{knot}}(B), HC_0^{\text{knot}}(\tilde{B})$ are isomorphic. By Markov's Theorem, it suffices to check this when B, \tilde{B} are related by one of three operations: conjugation, positive stabilization, and negative stabilization.

CONJUGATION. Here $B, \tilde{B} \in B_n$ satisfy $\tilde{B} = C^{-1}BC$ for some $C \in B_n$.

Write $HC_0^{\text{knot}}(B) = \mathcal{A}_n/\mathcal{I}$, where \mathcal{I} is generated by the entries of $(1 - \Phi_B^L(A)) \cdot A$ and $A \cdot (1 - \Phi_B^R(A))$. For clarity, write $HC_0^{\text{knot}}(\tilde{B}) = \tilde{\mathcal{A}}_n/\tilde{\mathcal{I}}$, with

$\tilde{\mathcal{A}}_n$ generated by $\{\tilde{a}_{ij}\}$ and $\tilde{\mathcal{I}}$ generated by the entries of $(1 + \Phi_B^L(\tilde{A})) \cdot \tilde{A}$ and $\tilde{A} \cdot (1 + \Phi_B^R(\tilde{A}))$, where $\tilde{A} = (\tilde{a}_{ij})$ and $\tilde{a}_{ii} = -2$.

We claim that the identification $\tilde{A} = \phi_C(A)$, i.e., $\tilde{a}_{ij} = \phi_C(a_{ij})$ for all i, j , maps $\tilde{\mathcal{I}}$ into \mathcal{I} . This would suffice to prove that $HC_0^{\text{knot}}(B) \cong HC_0^{\text{knot}}(\tilde{B})$, since a symmetrical argument with B and \tilde{B} interchanged shows that the same identification in the other direction also maps \mathcal{I} into $\tilde{\mathcal{I}}$, and hence \mathcal{I} and $\tilde{\mathcal{I}}$ coincide.

Assume that $\tilde{A} = \phi_C(A)$. From Proposition 4.4 and Corollary 4.5, we have

$$\Phi_B^L(\tilde{A}) = \Phi_{BC}^L(A) \cdot \Phi_{C^{-1}}^L(\phi_C(A)) = \Phi_C^L(\phi_B(A)) \cdot \Phi_B^L(A) \cdot (\Phi_C^L(A))^{-1},$$

and similarly $\Phi_B^R(\tilde{A}) = (\Phi_C^R(A))^{-1} \cdot \Phi_B^R(A) \cdot \Phi_C^R(\phi_B(A))$. It follows by Proposition 4.7 that

$$\begin{aligned} (1 - \Phi_B^L(\tilde{A})) \cdot \tilde{A} &= (1 - \Phi_C^L(\phi_B(A)) \cdot \Phi_B^L(A) \cdot (\Phi_C^L(A))^{-1}) \cdot \Phi_C^L(A) \cdot A \cdot \Phi_C^R(A) \\ &= \Phi_C^L(A) \cdot (1 - \Phi_B^L(A)) \cdot A \cdot \Phi_C^R(A) \\ &\quad - (\Phi_C^L(\phi_B(A)) - \Phi_C^L(A)) \cdot \Phi_B^L(A) \cdot A \cdot \Phi_C^R(A). \end{aligned}$$

Now the entries of $(1 - \Phi_B^L(A)) \cdot A$ are in \mathcal{I} by definition; on the other hand, the entries of

$$\phi_B(A) - A = \Phi_B^L(A) \cdot A \cdot \Phi_B^R(A) - A = -(1 - \Phi_B^L(A)) \cdot A \cdot \Phi_B^R(A) - A \cdot (1 - \Phi_B^R(A))$$

are in \mathcal{I} , and thus so are the entries of $\Phi_C^L(\phi_B(A)) - \Phi_C^L(A)$. Hence the entries of $(1 - \Phi_B^L(\tilde{A})) \cdot \tilde{A}$ are in \mathcal{I} . Similarly, the entries of $\tilde{A} \cdot (1 - \Phi_B^R(\tilde{A}))$ are also in \mathcal{I} , and therefore $\tilde{\mathcal{I}} \subset \mathcal{I}$, as desired.

POSITIVE STABILIZATION. Here \tilde{B} is obtained from B by adding an extra strand, which we label 0, to the braid, and setting $\tilde{B} = B\sigma_0$.

Analogously to before, write $HC_0^{\text{knot}}(B) = \mathcal{A}_n/\mathcal{I}$ and $HC_0^{\text{knot}}(\tilde{B}) = \tilde{\mathcal{A}}/\tilde{\mathcal{I}}$, where now $\tilde{\mathcal{A}}$ is obtained from \mathcal{A}_n by adding generators a_{i0}, a_{0i} for $1 \leq i \leq n$. Abbreviate $\Phi_B^L(A), \Phi_B^R(A)$ by Φ^L, Φ^R . By using Proposition 4.4 and Lemma 4.6, we can easily compute the matrices for $\Phi_{B\sigma_0}^L, \Phi_{B\sigma_0}^R$ in terms of Φ^L, Φ^R :

$$\begin{aligned} (\Phi_{B\sigma_0}^L)_{00} &= -\phi_B(a_{10}) & (\Phi_{B\sigma_0}^R)_{00} &= -\phi_B(a_{01}) \\ (\Phi_{B\sigma_0}^L)_{0i} &= -\Phi_{1i}^L & (\Phi_{B\sigma_0}^R)_{i0} &= -\Phi_{i1}^R \\ (\Phi_{B\sigma_0}^L)_{10} &= 1 & (\Phi_{B\sigma_0}^R)_{01} &= 1 \\ (\Phi_{B\sigma_0}^L)_{1i} &= 0 & (\Phi_{B\sigma_0}^R)_{i1} &= 0 \\ (\Phi_{B\sigma_0}^L)_{j0} &= 0 & (\Phi_{B\sigma_0}^R)_{0j} &= 0 \\ (\Phi_{B\sigma_0}^L)_{ji} &= \Phi_{ji}^L & (\Phi_{B\sigma_0}^R)_{ij} &= \Phi_{ij}^R. \end{aligned}$$

Here i, j are any indices such that $i \geq 1$ and $j \geq 2$. It is then straightforward to calculate the generators of $\tilde{\mathcal{I}}$; they are the entries of $(1 - \Phi_B^L(A)) \cdot A$ and

$A \cdot (1 - \Phi_B^R(A))$, which we write as $\tilde{\partial}\tilde{B}$ and $\tilde{\partial}\tilde{C}$ for future use in Section 5.2:

$$\begin{aligned}
\tilde{\partial}\tilde{b}_{00} &= -2 - \phi_B(a_{10}) & \tilde{\partial}\tilde{c}_{00} &= -2 - \phi_B(a_{01}) \\
\tilde{\partial}\tilde{b}_{0i} &= a_{0i} + \phi_B(a_{10})a_{0i} + \Phi_{1\ell}^L a_{\ell i} & \tilde{\partial}\tilde{c}_{i0} &= a_{i0} + a_{i0}\phi_B(a_{01}) + a_{i\ell}\Phi_{\ell 1}^R \\
\tilde{\partial}\tilde{b}_{10} &= a_{10} + 2 & \tilde{\partial}\tilde{c}_{01} &= a_{01} + 2 \\
\tilde{\partial}\tilde{b}_{1i} &= a_{1i} - a_{0i} & \tilde{\partial}\tilde{c}_{i1} &= a_{i1} - a_{i0} \\
\tilde{\partial}\tilde{b}_{j0} &= a_{j0} - \Phi_{j\ell}^L a_{\ell 0} & \tilde{\partial}\tilde{c}_{0j} &= a_{0j} - a_{0\ell}\Phi_{\ell j}^R \\
\tilde{\partial}\tilde{b}_{ji} &= a_{ji} - \Phi_{j\ell}^L a_{\ell i} & \tilde{\partial}\tilde{c}_{ij} &= a_{ij} - a_{i\ell}\Phi_{\ell j}^R,
\end{aligned}$$

where, as before, $i \geq 1$ and $j \geq 2$, and any monomial involving the index ℓ is understood to be summed from $\ell = 1$ to $\ell = n$. (To obtain the expressions for \tilde{b}_{00} and \tilde{c}_{00} , we use the facts that $\phi_B(a_{10}) = \Phi_{1\ell}^L a_{\ell 0}$ and $\phi_B(a_{01}) = a_{0\ell}\Phi_{\ell 1}^R$, which follow from the definitions of Φ^L and Φ^R .)

In $\tilde{\mathcal{A}}/\tilde{\mathcal{I}}$, we thus have $a_{0i} = a_{1i}$ and $a_{i0} = a_{i1}$ for $i \geq 1$; in particular, $a_{01} = a_{10} = -2$. Using these relations to replace all generators of the form a_{0i} and a_{i0} , we find that the remaining relations in $\tilde{\mathcal{A}}/\tilde{\mathcal{I}}$ give precisely the generators of \mathcal{I} , namely $a_{ij} - \Phi_{i\ell}^L a_{\ell j}$ and $a_{ij} - a_{i\ell}\Phi_{\ell j}^R$ for $1 \leq i, j \leq n$. Thus $\mathcal{A}_n/\mathcal{I} \cong \tilde{\mathcal{A}}/\tilde{\mathcal{I}}$, as desired.

NEGATIVE STABILIZATION. Here \tilde{B} is obtained from B by adding an extra strand, which we label 0, to the braid, and setting $\tilde{B} = B\sigma_0^{-1}$.

This case is very similar to the case of positive stabilization, with a slightly different computation. Use the same notation as for positive stabilization; then we have

$$\begin{aligned}
(\Phi_{B\sigma_0}^L)_{00} &= 0 & (\Phi_{B\sigma_0}^R)_{00} &= 0 \\
(\Phi_{B\sigma_0}^L)_{0i} &= \Phi_{1i}^L & (\Phi_{B\sigma_0}^R)_{i0} &= \Phi_{i1}^R \\
(\Phi_{B\sigma_0}^L)_{10} &= -1 & (\Phi_{B\sigma_0}^R)_{01} &= -1 \\
(\Phi_{B\sigma_0}^L)_{1i} &= -\phi_B(a_{01})\Phi_{1i}^L & (\Phi_{B\sigma_0}^R)_{i1} &= -\Phi_{i1}^R\phi_B(a_{10}) \\
(\Phi_{B\sigma_0}^L)_{j0} &= 0 & (\Phi_{B\sigma_0}^R)_{0j} &= 0 \\
(\Phi_{B\sigma_0}^L)_{ji} &= \Phi_{ji}^L & (\Phi_{B\sigma_0}^R)_{ij} &= \Phi_{ij}^R
\end{aligned}$$

for $i \geq 1$ and $j \geq 2$, and so the generators of $\tilde{\mathcal{I}}$ are

$$\begin{aligned}
\tilde{\partial}\tilde{b}_{00} &= -2 - \phi_B(a_{10}) & \tilde{\partial}\tilde{c}_{00} &= -2 - \phi_B(a_{01}) \\
\tilde{\partial}\tilde{b}_{0i} &= a_{0i} - \Phi_{1\ell}^L a_{\ell i} & \tilde{\partial}\tilde{c}_{i0} &= a_{i0} - a_{i\ell}\Phi_{\ell 1}^R \\
\tilde{\partial}\tilde{b}_{10} &= a_{10} - 2 + \phi_B(a_{01})\phi_B(a_{10}) & \tilde{\partial}\tilde{c}_{01} &= a_{01} - 2 + \phi_B(a_{01})\phi_B(a_{10}) \\
\tilde{\partial}\tilde{b}_{1i} &= a_{1i} + a_{0i} + \phi_B(a_{01})\Phi_{1\ell}^L a_{\ell i} & \tilde{\partial}\tilde{c}_{i1} &= a_{i1} + a_{i0} + a_{i\ell}\Phi_{\ell 1}^R\phi_B(a_{10}) \\
\tilde{\partial}\tilde{b}_{j0} &= a_{j0} - \Phi_{j\ell}^L a_{\ell 0} & \tilde{\partial}\tilde{c}_{0j} &= a_{0j} - a_{0\ell}\Phi_{\ell j}^R \\
\tilde{\partial}\tilde{b}_{ji} &= a_{ji} - \Phi_{j\ell}^L a_{\ell i} & \tilde{\partial}\tilde{c}_{ij} &= a_{ij} - a_{i\ell}\Phi_{\ell j}^R.
\end{aligned}$$

In $\tilde{\mathcal{A}}/\tilde{\mathcal{I}}$, we have $a_{0i} = \Phi_{1\ell}^L a_{\ell i}$ and $a_{i0} = a_{i\ell} \Phi_{\ell 1}^R$. When we use these relations to eliminate a_{0i}, a_{i0} , we find that the remaining relations in $\tilde{\mathcal{A}}/\tilde{\mathcal{I}}$ are precisely the relations in $\mathcal{A}_n/\mathcal{I}$, as before. Hence $\tilde{\mathcal{A}}/\tilde{\mathcal{I}} \cong \mathcal{A}_n/\mathcal{I}$.

We have shown that $HC_0^{\text{knot}}(B)$ is invariant under the Markov moves, and hence it gives a well-defined knot invariant, as desired. \square

5. INVARIANCE PROOFS FOR BRAID AND KNOT DGAS

In this section, we prove Theorems 2.7 and 2.10, the invariance results for braid and knot DGAs. As mentioned before, the proofs are essentially more involved versions of the corresponding proofs for HC_0 given in Section 4.

5.1. Proof of Theorem 2.7. It suffices to establish Theorem 2.7 under the assumption that $\tilde{B} = \sigma_k^{-1} B \sigma_k$ for some k . Then the braid DGAs for B and \tilde{B} are generated by a_{ij}, b_{ij} and $\tilde{a}_{ij}, \tilde{b}_{ij}$, respectively, for $1 \leq i \neq j \leq n$. If we abbreviate ϕ_B by ϕ and ϕ_{σ_k} by ϕ_k , then the differentials for the braid DGAs are given by $\partial b_{ij} = a_{ij} - \phi(a_{ij})$ and $\tilde{\partial} \tilde{b}_{ij} = \tilde{a}_{ij} - \phi_k^{-1} \phi \phi_k(\tilde{a}_{ij})$. (By abusing notation, we will consider ϕ and ϕ_k to act either on the algebra generated by the a_{ij} or on the algebra generated by the \tilde{a}_{ij} .) We will find a tame isomorphism between the algebras which sends one differential to the other.

Set $\tilde{a}_{ij} = \phi_k(a_{ij})$ for all i, j . Then

$$\tilde{\partial} \tilde{b}_{ij} = \tilde{a}_{ij} - \phi_k^{-1} \phi \phi_k(\tilde{a}_{ij}) = \phi_k(a_{ij}) - \phi \phi_k(a_{ij}).$$

Now make the following identifications:

$$\begin{aligned} \tilde{b}_{ki} &= -b_{k+1,i} - \phi(a_{k+1,k}) b_{ki} - b_{k+1,k} a_{ki} & i \neq k, k+1 \\ \tilde{b}_{ik} &= -b_{i,k+1} - b_{ik} \phi(a_{k,k+1}) - a_{ik} b_{k,k+1} & i \neq k, k+1 \\ \tilde{b}_{k+1,i} &= b_{ki} & i \neq k, k+1 \\ \tilde{b}_{i,k+1} &= b_{ik} & i \neq k, k+1 \\ \tilde{b}_{k,k+1} &= b_{k+1,k} \\ \tilde{b}_{k+1,k} &= b_{k,k+1} \\ \tilde{b}_{ij} &= b_{ij} & i, j \neq k, k+1. \end{aligned}$$

It is straightforward to check from $\partial b_{ij} = a_{ij} - \phi(a_{ij})$ that $\tilde{\partial} \tilde{b}_{ij} = \phi_k(a_{ij}) - \phi \phi_k(a_{ij})$ for all i, j . For instance, for $i \neq k, k+1$, we have

$$\begin{aligned} \tilde{\partial} \tilde{b}_{ik} &= -\partial b_{i,k+1} - (\partial b_{ik}) \phi(a_{k,k+1}) - a_{ik} (\partial b_{k,k+1}) \\ &= -a_{i,k+1} + \phi(a_{i,k+1}) + \phi(a_{ik}) \phi(a_{k,k+1}) - a_{ik} a_{k,k+1} \\ &= \phi_k(a_{ik}) - \phi \phi_k(a_{ik}). \end{aligned}$$

Using the above definitions for $\tilde{a}_{ij}, \tilde{b}_{ij}$ in terms of a_{ij}, b_{ij} , we conclude that $\partial = \tilde{\partial}$, and so the map sending a_{ij} to \tilde{a}_{ij} and b_{ij} to \tilde{b}_{ij} sends ∂ to $\tilde{\partial}$. On the other hand, it is easy to check that this map is tame; the map on the

a_{ij} is given by the tame isomorphism ϕ_k , while the map on the b_{ij} is tame by inspection of the above definition for \tilde{b}_{ij} in terms of b_{ij} . Thus the braid DGAs for B and $\sigma_k^{-1}B\sigma_k$ are tamely isomorphic, as desired. \square

5.2. Proof of Theorem 2.10. As in the proof of Theorem 4.10, it suffices to show that the equivalence class of the knot DGA of a braid is invariant under the Markov moves.

CONJUGATION. Let $B \in B_n$ be a braid, and let $\tilde{B} = \sigma_k^{-1}B\sigma_k$; we show that the modified knot DGAs (see Proposition 4.8) for B and \tilde{B} are tamely isomorphic.

For clarity, we distinguish between the modified knot DGAs for B and \tilde{B} by using tildes on the generators and differential of the knot DGA for \tilde{B} . As in the proof of Theorem 2.7, we will exhibit an identification between the two sets of generators so that $\partial = \tilde{\partial}$.

By Lemma 4.6, the matrices $\Phi_{\sigma_k}^L(A) - \Phi_{\sigma_k}^L(\phi_B(A))$ and $\Phi_{\sigma_k}^R(A) - \Phi_{\sigma_k}^R(\phi_B(A))$ are both identically zero except in the (k, k) entry, where they are $-a_{k+1,k} + \phi_B(a_{k+1,k})$ and $-a_{k,k+1} + \phi_B(a_{k,k+1})$, respectively. Hence if we define $n \times n$ matrices $\Theta_k^L(A)$ and $\Theta_k^R(A)$ to be zero except in the (k, k) entry, where they are $-b_{k+1,k}$ and $-b_{k,k+1}$, respectively, then we have

$$\partial\Theta_k^L(A) = \Phi_{\sigma_k}^L(A) - \Phi_{\sigma_k}^L(\phi_B(A)) \quad \text{and} \quad \partial\Theta_k^R(A) = \Phi_{\sigma_k}^R(A) - \Phi_{\sigma_k}^R(\phi_B(A)).$$

Identify the $\tilde{a}_{ij}, \tilde{b}_{ij}$ with the a_{ij}, b_{ij} using the same map as in the proof of Theorem 2.7 in Section 5.1; then ∂ and $\tilde{\partial}$ agree on \tilde{a}_{ij} and \tilde{b}_{ij} . Now set

$$\begin{aligned} \tilde{C} &= \Phi_{\sigma_k}^L(\phi_B(A)) \cdot C \cdot \Phi_{\sigma_k}^R(A) + \Theta_k^L(A) \cdot A \cdot \Phi_{\sigma_k}^R(A), \\ \tilde{D} &= \Phi_{\sigma_k}^L(A) \cdot D \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \Phi_{\sigma_k}^L(A) \cdot A \cdot \Theta_k^R(A), \\ \tilde{E} &= \Phi_{\sigma_k}^L(\phi_B(A)) \cdot E \cdot \Phi_{\sigma_k}^R(\phi_B(A)) - \Theta_k^L(A) \cdot D \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \Theta_k^L(A) \cdot \Theta_k^R(A), \\ \tilde{F} &= \Phi_{\sigma_k}^L(\phi_B(A)) \cdot F \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \Phi_{\sigma_k}^L(\phi_B(A)) \cdot C \cdot \Theta_k^R(A) + \Theta_k^L(A) \cdot (1 + A) \cdot \Theta_k^R(A). \end{aligned}$$

We claim that these identifications make ∂ and $\tilde{\partial}$ agree on $\tilde{c}_{ij}, \tilde{d}_{ij}, \tilde{e}_{ij}, \tilde{f}_{ij}$, which implies that $\partial = \tilde{\partial}$ on the entire algebra. We will check that $\tilde{\partial}\tilde{C} = \partial\tilde{C}$ and $\tilde{\partial}\tilde{E} = \partial\tilde{E}$; the proofs for D and F are completely analogous.

By Proposition 4.4 and Corollary 4.5, we have

$$\Phi_{\tilde{B}}^L(\tilde{A}) = \Phi_{\sigma_k^{-1}B\sigma_k}^L(\phi_{\sigma_k}(A)) = \Phi_{B\sigma_k}^L(A) \cdot \Phi_{\sigma_k^{-1}}^L(\phi_{\sigma_k}(A)) = \Phi_{\sigma_k}^L(\phi_B(A)) \cdot \Phi_B^L(A) \cdot (\Phi_{\sigma_k}^L(A))^{-1},$$

and similarly

$$\Phi_{\tilde{B}}^R(\tilde{A}) = (\Phi_{\sigma_k}^R(A))^{-1} \cdot \Phi_B^R(A) \cdot \Phi_{\sigma_k}^R(\phi_B(A)).$$

Recall from Proposition 4.7 that $\tilde{A} = \phi_{\sigma_k}(A) = \Phi_{\sigma_k}^L(A) \cdot A \cdot \Phi_{\sigma_k}^R(A)$; it follows that

$$\begin{aligned} \partial \tilde{C} &= \Phi_{\sigma_k}^L(\phi_B(A)) \cdot (1 - \Phi_B^L(A)) \cdot A \cdot \Phi_{\sigma_k}^R(A) + (\Phi_{\sigma_k}^L(A) - \Phi_{\sigma_k}^L(\phi_B(A))) \cdot A \cdot \Phi_{\sigma_k}^R(A) \\ &= (\Phi_{\sigma_k}^L(A) - \Phi_{\sigma_k}^L(\phi_B(A)) \cdot \Phi_B^L(A)) \cdot A \cdot \Phi_{\sigma_k}^R(A) \\ &= (1 - \Phi_B^L(\tilde{A})) \cdot \tilde{A} \\ &= \tilde{\partial} \tilde{C} \end{aligned}$$

and

$$\begin{aligned} \partial \tilde{E} &= \Phi_{\sigma_k}^L(\phi_B(A)) \cdot (B - D - C \cdot \Phi_B^R(A)) \cdot \Phi_{\sigma_k}^R(\phi_B(A)) \\ &\quad - (\Phi_{\sigma_k}^L(A) - \Phi_{\sigma_k}^L(\phi_B(A))) \cdot D \cdot \Phi_{\sigma_k}^R(\phi_B(A)) \\ &\quad + \Theta_k^L(A) \cdot A \cdot (1 - \Phi_B^R(A)) \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \partial(\Theta_k^L(A) \cdot \Theta_k^R(A)) \\ &= \tilde{B} - \tilde{D} - \tilde{C} \cdot \Phi_{\tilde{B}}^R(\tilde{A}) \\ &= \tilde{\partial} \tilde{E}, \end{aligned}$$

where we use the fact (a direct computation from the definition of \tilde{b}_{ij}) that

$$\begin{aligned} \tilde{B} &= \Phi_{\sigma_k}^L(\phi_B(A)) \cdot B \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \Phi_{\sigma_k}^L(A) \cdot A \cdot \Theta_k^R(A) \\ &\quad + \Theta_k^L(A) \cdot A \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \partial(\Theta_k^L(A) \cdot \Theta_k^R(A)). \end{aligned}$$

Now that we have established that $\partial = \tilde{\partial}$ on the entire algebra, it only remains to verify that the identification of variables $A \mapsto \tilde{A}$, etc., constitutes a tame isomorphism. This is straightforward; as in the proof of Theorem 2.7, the map $A \mapsto \tilde{A}, B \mapsto \tilde{B}$ is a tame automorphism. We can then send C to \tilde{C} via the composition of tame automorphisms

$$\begin{aligned} C &\mapsto C + \Theta_k^L(A) \cdot A \cdot \Phi_{\sigma_k}^R(A), \\ C &\mapsto \Phi_{\sigma_k}^L(\phi_B(A)) \cdot C, \\ C &\mapsto C \cdot \Phi_{\sigma_k}^R(A), \end{aligned}$$

of which the last two are tame because of Lemma 4.6, and similarly for $D \mapsto \tilde{D}$. Finally, we can send E to \tilde{E} via the composition of tame automorphisms

$$\begin{aligned} E &\mapsto E - \Theta_k^L(A) \cdot D \cdot \Phi_{\sigma_k}^R(\phi_B(A)) + \Theta_k^L(A) \cdot \Theta_k^R(A), \\ E &\mapsto \Phi_{\sigma_k}^L(\phi_B(A)) \cdot E, \\ E &\mapsto E \cdot \Phi_{\sigma_k}^R(\phi_B(A)), \end{aligned}$$

and similarly for $F \mapsto \tilde{F}$.

POSITIVE STABILIZATION. Let $B \in B_n$ be a braid; as in the proof of Theorem 4.10, we denote stabilization by adding a strand labelled 0 and considering the braid $\tilde{B} = B\sigma_0$. We will prove that the knot DGAs for B and \tilde{B} are stable tame isomorphic.

Denote the knot DGA for B by (\mathcal{A}, ∂) and the knot DGA for \tilde{B} by $(\tilde{\mathcal{A}}, \tilde{\partial})$, where the generators of $\tilde{\mathcal{A}}$ have tildes for notational clarity. Thus $\tilde{\mathcal{A}}$ is

generated by $\{\tilde{a}_{ij} \mid 0 \leq i \neq j \leq n\}$ in degree 0, and so forth. Note that $\tilde{\mathcal{A}}$ has more generators than \mathcal{A} does; we will establish that we can obtain $(\tilde{\mathcal{A}}, \tilde{\partial})$ from (\mathcal{A}, ∂) by a suitable number of stabilizations.

We will progressively identify generators of \mathcal{A} and $\tilde{\mathcal{A}}$ so that ∂ and $\tilde{\partial}$ agree. The first identification is to drop all tildes on the \tilde{a}_{ij} generators; that is, set $a_{ij} = \tilde{a}_{ij}$ for all $i, j \geq 0, i \neq j$. Note that this introduces $2n$ generators a_{0j} and a_{i0} not in \mathcal{A} .

The matrices $\Phi_{B\sigma_0}^L$ and $\Phi_{B\sigma_0}^R$ were computed in the proof of Theorem 4.10. We can then compute $\tilde{\partial}$ in terms of $\Phi^L = \Phi_B^L$ and $\Phi^R = \Phi_B^R$. For \tilde{b}_{ij} and \tilde{c}_{ij} , we already calculated $\tilde{\partial}$ in the proof of Theorem 4.10. For completeness, we include here the differentials of \tilde{d}_{ij} and \tilde{e}_{ij} :

$$\begin{aligned}
\tilde{\partial}\tilde{d}_{00} &= \tilde{b}_{00} - \tilde{c}_{00} + \tilde{b}_{00}\phi_B(a_{01}) + \tilde{b}_{0\ell}\Phi_{\ell 1}^R - \phi_B(a_{10})\tilde{c}_{00} - \Phi_{1\ell}^L\tilde{c}_{\ell 0} \\
\tilde{\partial}\tilde{d}_{01} &= \tilde{b}_{01} - \tilde{c}_{01} - \tilde{b}_{00} - \phi_B(a_{10})\tilde{c}_{01} - \Phi_{1\ell}^L\tilde{c}_{\ell 1} \\
\tilde{\partial}\tilde{d}_{0j} &= \tilde{b}_{0j} - \tilde{c}_{0j} - \tilde{b}_{0\ell}\Phi_{\ell j}^R - \phi_B(a_{10})\tilde{c}_{0j} - \Phi_{1\ell}^L\tilde{c}_{\ell j} \\
\tilde{\partial}\tilde{d}_{10} &= \tilde{b}_{10} - \tilde{c}_{10} + \tilde{c}_{00} + \tilde{b}_{10}\phi_B(a_{01}) + \tilde{b}_{1\ell}\Phi_{\ell 1}^R \\
\tilde{\partial}\tilde{d}_{11} &= \tilde{b}_{11} - \tilde{c}_{11} - \tilde{b}_{10} + \tilde{c}_{01} \\
\tilde{\partial}\tilde{d}_{1j} &= \tilde{b}_{1j} - \tilde{c}_{1j} + \tilde{c}_{0j} - \tilde{b}_{1\ell}\Phi_{\ell 1}^R \\
\tilde{\partial}\tilde{d}_{j0} &= \tilde{b}_{j0} - \tilde{c}_{j0} + \tilde{b}_{j0}\phi_B(a_{01}) + \tilde{b}_{j\ell}\Phi_{\ell 1}^R + \Phi_{j\ell}^L\tilde{c}_{\ell 0} \\
\tilde{\partial}\tilde{d}_{j1} &= \tilde{b}_{j1} - \tilde{c}_{j1} - \tilde{b}_{j0} + \Phi_{j\ell}^L\tilde{c}_{\ell 1} \\
\tilde{\partial}\tilde{d}_{j_1j_2} &= \tilde{b}_{j_1j_2} - \tilde{c}_{j_1j_2} - \tilde{b}_{j_1\ell}\Phi_{\ell j_2}^R + \Phi_{j_1\ell}^L\tilde{c}_{\ell j_2} \\
\tilde{\partial}\tilde{e}_0 &= \tilde{b}_{00} - \phi_B(a_{10})\tilde{c}_{00} - \Phi_{1\ell}^L\tilde{c}_{\ell 0} \\
\tilde{\partial}\tilde{e}_1 &= \tilde{b}_{11} + \tilde{c}_{01} \\
\tilde{\partial}\tilde{e}_j &= \tilde{b}_{jj} + \Phi_{j\ell}^L\tilde{c}_{\ell j},
\end{aligned}$$

where $j, j_1, j_2 \geq 2$ and, as usual, all monomials involving ℓ are summed from $\ell = 1$ to $\ell = n$.

If we set

$$\begin{aligned}
b_{1i} &= -\tilde{b}_{0i} + \tilde{b}_{1i} - \tilde{b}_{00}a_{0i} & c_{i1} &= -\tilde{c}_{i0} + \tilde{c}_{i1} - a_{i0}\tilde{c}_{00} \\
b_{ji} &= \tilde{b}_{ji} & c_{ij} &= \tilde{c}_{ij}
\end{aligned}$$

for $i \geq 1$ and $j \geq 2$, then it is easy to check, from the expressions for $\tilde{\partial}\tilde{b}_{ij}$, $\tilde{\partial}\tilde{c}_{ij}$ given in the proof of Theorem 4.10, that $\tilde{\partial}b_{ij} = \partial b_{ij}$ and $\tilde{\partial}c_{ij} = \partial c_{ij}$ for

all $i, j \geq 1$. Next set

$$\begin{aligned}
 d_{11} &= \tilde{d}_{00} - \tilde{d}_{01} + \tilde{d}_{11} - \tilde{d}_{10} + \tilde{b}_{00}\tilde{c}_{01} + \tilde{b}_{10}\tilde{c}_{00} \\
 d_{1j} &= -\tilde{d}_{0j} + \tilde{d}_{1j} + \tilde{b}_{00}\tilde{c}_{0j} \\
 d_{j1} &= -\tilde{d}_{j0} + \tilde{d}_{j1} + \tilde{b}_{j0}\tilde{c}_{00} \\
 d_{j_1j_2} &= \tilde{d}_{j_1j_2} \\
 e_1 &= \tilde{e}_0 + \tilde{e}_1 - \tilde{d}_{01} + \tilde{b}_{00}\tilde{c}_{01} \\
 e_j &= \tilde{e}_j \\
 d_{00} &= \tilde{d}_{10} - \tilde{d}_{11} + \tilde{d}_{01} - \tilde{b}_{10}\tilde{c}_{00} - \tilde{b}_{00}\tilde{c}_{01} \\
 e_0 &= \tilde{d}_{01} - \tilde{e}_1 - \tilde{b}_{00}\tilde{c}_{01} \\
 d_{10} &= \tilde{d}_{11} - \tilde{e}_1 & d_{01} &= \tilde{e}_1 & d_{0j} &= \tilde{d}_{1j} & d_{j0} &= \tilde{d}_{j1}
 \end{aligned}$$

for $j, j_1, j_2 \geq 2$, and then successively set

$$\begin{aligned}
 b_{10} &= \tilde{b}_{10} + \tilde{c}_{11} & \text{and} & & c_{01} &= \tilde{c}_{01} + \tilde{b}_{11}; \\
 b_{00} &= \tilde{b}_{00} - b_{11} - \Phi_{1\ell}^L \tilde{c}_{\ell 1} & \text{and} & & c_{00} &= \tilde{c}_{00} - c_{11} - \tilde{b}_{1\ell} \Phi_{\ell 1}^R; \\
 b_{j0} &= \tilde{b}_{j0} + \tilde{c}_{j1} - b_{j1} - \Phi_{j\ell}^L \tilde{c}_{\ell 1} & \text{and} & & c_{0j} &= \tilde{c}_{0j} + \tilde{b}_{1j} - c_{1j} - \tilde{b}_{1\ell} \Phi_{\ell j}^R; \\
 b_{0i} &= \tilde{b}_{1i} & \text{and} & & c_{i0} &= \tilde{c}_{i1}
 \end{aligned}$$

for $i \geq 1$ and $j \geq 2$.

Under the above identifications, the map sending the tilde variables to the corresponding non-tilde variables is a tame automorphism on $\tilde{\mathcal{A}}$. To see this, start with the generators of $\tilde{\mathcal{A}}$, and now perform the following changes of variables. First introduce new variables b_{1i}, c_{i1} for $i \geq 1$, and then eliminate the generators $\tilde{b}_{0i}, \tilde{c}_{i0}$ by using the first identifications: $\tilde{b}_{0i} = -b_{1i} + \tilde{b}_{1i} - \tilde{b}_{00}a_{0i}$ and $\tilde{c}_{i0} = -c_{i1} + \tilde{c}_{i1} - a_{i0}\tilde{c}_{00}$. Now successively continue this process using the given identifications, from first to last; for each identification, introduce a new generator given by the left hand side, and use the relation to eliminate the generator given by the first term on the right hand side. This can be done because no eliminated generator appears in successive identifications. The final result is that all tilde variables are eliminated and replaced by non-tilde variables. It follows that the map sending tilde generators to non-tilde generators is tame.

On the other hand, a tedious but straightforward series of calculations demonstrates that $\partial = \tilde{\partial}$ after the above identifications have been made, where we set $\partial b_{0i} = -a_{0i} + a_{1i}$, $\partial c_{i0} = -a_{i0} + a_{i1}$, $\partial e_0 = b_{00}$, $\partial d_{00} = b_{00} - c_{00}$, $\partial d_{i0} = -b_{i0}$, $\partial d_{0i} = c_{0i}$ for $i \geq 1$. It follows that $(\tilde{\mathcal{A}}, \tilde{\partial})$ is tamely isomorphic to the result of stabilizing (\mathcal{A}, ∂) by appending generators $a_{0i}, a_{i0}, b_{00}, b_{0i}, b_{i0}, c_{00}, c_{0i}, c_{i0}, d_{00}, d_{0i}, d_{i0}, e_0$, and extending ∂ as in the previous sentence. (This corresponds to $2n$ stabilizations of degree 0 and $2n + 2$ stabilizations of degree 1.) This completes the proof that the knot DGAs for B and \tilde{B} are stable tame isomorphic.

NEGATIVE STABILIZATION. We now wish to prove that the knot DGAs for B and $\tilde{B} = B\sigma_0^{-1}$ are stable tame isomorphic; this proof is entirely similar to the proof for positive stabilization. We omit the calculations, but the sequence of successive identifications is as follows, where indices satisfy $i \geq 1$ and $j, j_1, j_2 \geq 2$, and all monomials involving ℓ are understood to be summed over $\ell \geq 1$:

$$\begin{aligned} b_{1i} &= \tilde{b}_{1i} - \tilde{b}_{0i} + \tilde{c}_{00}\Phi_{1\ell}^L a_{\ell i} & c_{i1} &= \tilde{c}_{i1} - \tilde{c}_{i0} + a_{i\ell}\Phi_{\ell 1}^R \tilde{b}_{00} \\ b_{ji} &= \tilde{b}_{ji} & c_{ij} &= \tilde{c}_{ij}, \end{aligned}$$

followed by

$$\begin{aligned} d_{11} &= -\tilde{d}_{10} + \tilde{d}_{11} - \tilde{d}_{01} + \tilde{d}_{00} - b_{1\ell}\Phi_{\ell 1}^R \tilde{b}_{00} - \tilde{c}_{00}\Phi_{1\ell}^L c_{\ell 1} - 2\tilde{c}_{00}\tilde{b}_{00} \\ d_{1j} &= \tilde{d}_{1j} - \tilde{d}_{0j} - \tilde{c}_{00}\Phi_{1\ell}^L c_{\ell j} \\ d_{j1} &= \tilde{d}_{j1} - \tilde{d}_{j0} - b_{j\ell}\Phi_{\ell 1}^R \tilde{b}_{00} \\ d_{j_1 j_2} &= \tilde{d}_{j_1 j_2} \\ e_1 &= -\tilde{d}_{01} + \tilde{e}_0 + \tilde{e}_1 - \tilde{c}_{00}\Phi_{1\ell}^L c_{\ell 1} - \tilde{c}_{00}\tilde{b}_{00} + (\tilde{e}_0 - \tilde{d}_{00})\phi_B(a_{10}) + \phi_B(a_{01})\tilde{e}_0 \\ e_j &= \tilde{e}_j \\ d_{00} &= \tilde{d}_{00} & e_0 &= \tilde{e}_0 & d_{10} &= \tilde{d}_{11} - \tilde{e}_1 & d_{01} &= \tilde{e}_1 & d_{j0} &= \tilde{d}_{j0} & d_{0j} &= \tilde{d}_{0j}, \end{aligned}$$

and finally

$$\begin{aligned} b_{10} &= -\tilde{b}_{10} + c_{11} + \tilde{c}_{10} - a_{1\ell}\Phi_{\ell 1}^R \tilde{b}_{00} & \text{and} & & c_{01} &= -\tilde{c}_{01} + b_{11} + \tilde{b}_{01} - \tilde{c}_{00}\Phi_{1\ell}^L a_{\ell 1} \\ & - (b_{1\ell}\Phi_{\ell 1}^R + \tilde{b}_{0\ell}\Phi_{\ell 1}^R + 2\tilde{c}_{00})\phi_B(a_{10}) & & & & - \phi_B(a_{01})(\Phi_{1\ell}^L c_{\ell 1} + \Phi_{1\ell}^L \tilde{c}_{\ell 0} + 2\tilde{b}_{00}); \\ b_{00} &= \tilde{b}_{00} + \Phi_{1\ell}^L \tilde{c}_{\ell 0} & \text{and} & & c_{00} &= \tilde{c}_{00} + \tilde{b}_{0\ell}\Phi_{\ell 1}^R; \\ b_{j0} &= -\tilde{b}_{j0} + \tilde{c}_{j0} + b_{j\ell}\Phi_{\ell 1}^R - \Phi_{1\ell}^L \tilde{c}_{\ell 0} & \text{and} & & c_{0j} &= -\tilde{c}_{0j} + \tilde{b}_{0j} + \Phi_{1\ell}^L c_{\ell j} - \tilde{b}_{0\ell}\Phi_{\ell j}^R; \\ b_{0i} &= \tilde{b}_{0i} & \text{and} & & c_{i0} &= \tilde{c}_{i0}. \end{aligned}$$

We find that $\partial = \tilde{\partial}$ as before, where we extend ∂ by defining $\partial b_{0i} = a_{0i} - \Phi_{1\ell}^L a_{\ell i}$, $\partial c_{i0} = a_{i0} - a_{i\ell}\Phi_{\ell 1}^R$, $\partial d_{00} = b_{00} - c_{00}$, $\partial e_0 = b_{00}$, $\partial d_{i0} = -b_{i0}$, $\partial d_{0i} = c_{0i}$. (The calculation uses the fact that $\Phi_{1\ell}^L a_{\ell m}\Phi_{m1}^R = (\phi_B(A))_{11} = -2$.) It follows that the knot DGA for $B\sigma_0^{-1}$ is tamely isomorphic to a stabilization of the knot DGA for B , as desired. \square

6. PROPERTIES OF THE INVARIANTS

6.1. Conjugation and abelianization. There is a symmetry in the homomorphism $\phi: B_n \rightarrow \text{Aut}(\mathcal{A}_n)$ which induces a symmetry in braid and knot DGAs. Define an involution on \mathcal{A}_n , which we term *conjugation* and write as $v \mapsto \bar{v}$, as follows: the conjugate of a monomial $a_{i_1 j_1} \cdots a_{i_m j_m}$ is $a_{j_m i_m} \cdots a_{j_1 i_1}$; extend conjugation linearly, and set $\bar{1} = 1$. In other words, conjugation replaces each generator a_{ij} by a_{ji} and then reverses the order of each word. The key observation is the following.

Proposition 6.1. *For all $v \in \mathcal{A}_n$ and any $B \in B_n$, we have $\phi_B(\bar{v}) = \overline{\phi_B(v)}$.*

Proof. The result holds for $B = \sigma_k$ and $v = a_{ij}$ for any i, j, k , by direct inspection of the definition of ϕ . The general result for $B = \sigma_k$ follows since $\overline{vw} = \overline{wv}$ for all $v, w \in \mathcal{A}_n$; since ϕ is a homomorphism, the entire proposition follows. \square

One consequence of Proposition 6.1 is that the matrices $\Phi_B^L(A), \Phi_B^R(A)$ determine each other. Extend the operation of conjugation to matrices by conjugating each entry, and let T denote transpose; then we have the following result.

Proposition 6.2. *For any $B \in B_n$, $\Phi_B^R(A) = \overline{\Phi_B^L(A)}^T$.*

Proof. By Proposition 6.1, we have

$$\phi_B(a_{*i}) = \phi_B(\overline{a_{i*}}) = \overline{\phi_B(a_{i*})} = \overline{\sum_{j=1}^n (\Phi_B^L(A))_{ij} a_{j*}} = \sum_{j=1}^n a_{*j} \overline{(\Phi_B^L(A))_{ij}};$$

now use the definition of $\Phi_B^R(A)$. \square

Conjugation also allows us to define simplified commutative versions of the braid and knot DGAs. Intuitively, we can quotient the braid and knot DGAs by conjugation and abelianize; the differential is well-defined on the resulting quotient algebras. For instance, on the braid DGA, if $\partial b_{ij} = a_{ij} - \phi_B(a_{ij})$, then $\partial b_{ji} = a_{ji} - \phi_B(a_{ji}) = \overline{a_{ij} - \phi_B(a_{ij})}$, and so ∂ is still well-defined if we mod out by $a_{ji} = a_{ij}$, $b_{ji} = b_{ij}$ and then abelianize.

Definition 6.3. Let $B \in B_n$ be a braid. Let \mathcal{A} be the graded sign-commutative algebra on $n(n-1)$ generators, $\{a_{ij} \mid 1 \leq i < j \leq n\}$ of degree 0 and $\{b_{ij} \mid 1 \leq i < j \leq n\}$ of degree 1. Define a differential ∂ on \mathcal{A} to be the usual differential on the braid DGA, where we set $a_{ji} = a_{ij}$ for $j > i$. Then we call (\mathcal{A}, ∂) the *abelian braid DGA* of B .

Here by “sign-commutative,” we mean that $vw = (-1)^{(\deg v)(\deg w)}wv$ for all $v, w \in \mathcal{A}$ of pure degree.

For the knot DGA, extend the conjugation operation to the entire algebra by defining $\overline{b_{ij}} = c_{ji}$, $\overline{c_{ij}} = b_{ji}$, $\overline{d_{ij}} = -d_{ji}$, $\overline{e_i} = e_i - d_{ii}$, and $\overline{vw} = (-1)^{(\deg v)(\deg w)}\overline{wv}$.

Proposition 6.4. *If (\mathcal{A}, ∂) is the knot DGA for some braid, then $\overline{\partial v} = \partial \overline{v}$ for all $v \in \mathcal{A}$.*

Proof. We calculate directly that

$$\begin{aligned}\overline{\partial b_{ij}} &= \overline{a_{ij}} - \sum_{\ell} \overline{a_{\ell j}} \overline{(\Phi_B^L(A))_{i\ell}} = \partial c_{ji}, \\ \overline{\partial c_{ij}} &= \overline{a_{ij}} - \sum_{\ell} \overline{(\Phi_B^R(A))_{\ell j}} \overline{a_{i\ell}} = \partial b_{ji}, \\ \overline{\partial d_{ij}} &= \overline{b_{ij}} - \overline{c_{ij}} + \sum_{\ell} \left(-\overline{(\Phi_B^R(A))_{\ell j}} \overline{b_{i\ell}} + \overline{c_{\ell j}} \overline{(\Phi_B^L(A))_{i\ell}} \right) = -\partial d_{ji}, \\ \overline{\partial e_i} &= \overline{b_{ii}} + \sum_{\ell} \overline{c_{\ell i}} \overline{(\Phi_B^L(A))_{i\ell}} = c_{ii} + b_{i\ell} (\Phi_B^R(A))_{\ell i} = \partial e_i - \partial d_{ii},\end{aligned}$$

and if $\overline{\partial v} = \partial \overline{v}$ and $\overline{\partial w} = \partial \overline{w}$, then

$$\overline{\partial(vw)} = (-1)^{(\deg v-1)(\deg w)} \overline{w} \overline{\partial v} + (-1)^{(\deg v)(\deg w)} \overline{\partial w} \overline{v} = \partial(\overline{vw}).$$

The proposition follows. \square

Since ∂ commutes with conjugation, we may mod out by conjugation and abelianize, and ∂ is well-defined on the resulting algebra.

Definition 6.5. Let $B \in B_n$. Let \mathcal{A} be the graded sign-commutative algebra with generators $\{a_{ij} \mid 1 \leq i < j \leq n\}$ of degree 0, $\{b_{ij} \mid 1 \leq i, j \leq n\}$ of degree 1, and $\{d_{ij} \mid 1 \leq i < j \leq n\}$ and $\{e_i \mid 1 \leq i \leq n\}$ of degree 2. Define ∂ on \mathcal{A} to be the usual differential on the knot DGA, where we set $a_{ji} = a_{ij}$ for $j > i$ and $c_{ij} = b_{ji}$ for all i, j . Then we call (\mathcal{A}, ∂) the *abelian knot DGA* of B .

The notion of stable tame isomorphism can be defined for polynomial algebras just as for tensor algebras, and with respect to this relation, abelian braid and knot DGAs give invariants.

Proposition 6.6. *Up to stable tame isomorphism, the abelian braid DGA of B is an invariant of the conjugacy class of B , and the abelian knot DGA of B is an invariant of the knot closure of B .*

Proof. Trace through the invariance proofs for the original knot and braid DGAs, and note that all of the tame isomorphisms and stabilizations used there are invariant under conjugation and hence well-defined in the commutative case. \square

As in the noncommutative case, our main interest in the abelian DGAs lies in the homology in degree 0, which we write as $HC_0^{\text{ab}}(B)$ in the braid case and $HC_0^{\text{ab}}(K)$ in the knot case.

Corollary 6.7. *Up to isomorphism, $HC_0^{\text{ab}}(B)$ is an invariant of the conjugacy class of B , and $HC_0^{\text{ab}}(K)$ is an invariant of the knot K .*

We can summarize the various flavors of HC_0 in a diagram, as follows. Suppose that braid B has closure K , and let $\text{ab}(HC_0(B))$, $\text{ab}(HC_0(K))$ represent the abelianizations of $HC_0(B)$, $HC_0(K)$. Then $HC_0(B)$, $HC_0(K)$ can

both be written as $\mathbb{Z}\langle\{a_{ij}\}\rangle$ modulo some ideal, and the ideal for $HC_0(K)$ contains the ideal for $HC_0(B)$ by Proposition 4.8. Also, $\text{ab}(HC_0(B)), HC_0^{\text{ab}}(B)$ are both $\mathbb{Z}[\{a_{ij}\}]$ modulo some ideal, and the ideal for $HC_0^{\text{ab}}(B)$ is generated by the ideal for $\text{ab}(HC_0(B))$ and the relations $\{a_{ij} - a_{ji}\}$. Thus we have the following commutative diagram of invariants:

$$\begin{array}{ccccc} HC_0(B) & \twoheadrightarrow & \text{ab}(HC_0(B)) & \twoheadrightarrow & HC_0^{\text{ab}}(B) \\ \downarrow & & \downarrow & & \downarrow \\ HC_0(K) & \twoheadrightarrow & \text{ab}(HC_0(K)) & \twoheadrightarrow & HC_0^{\text{ab}}(K). \end{array}$$

6.2. Mirrors and inverses. In this section, we show that the braid and knot DGAs do not change under mirroring and inversion. Recall that the *mirror* of a knot K is the knot \overline{K} obtained by reversing each of the crossings of the diagram of K , and similarly for the mirror \overline{B} of a braid B . In B_n , mirroring is the homomorphism sending σ_k to σ_k^{-1} for each k . It will be convenient for us to consider a related mirror operation on B_n , $B \mapsto B^*$, which is the homomorphism sending σ_k to σ_{n-k}^{-1} for each k . Recall also that the *inverse* of a knot is the knot with the opposite orientation. If K is the closure of $B \in B_n$, then the mirror \overline{K} is the closure of both \overline{B} and B^* , while the inverse of K is the closure of \overline{B}^{-1} .

Proposition 6.8. *The braid DGAs for a braid B , its mirrors \overline{B} and B^* , and its inverse B^{-1} are stable tame isomorphic.*

Proof. Let $B \in B_n$. We first show that B and B^{-1} have equivalent braid DGAs. In the braid DGA of B^{-1} , we have $\partial b_{ij} = a_{ij} - \phi_{B^{-1}}(a_{ij})$ for all i, j ; hence if we replace a_{ij} by $\phi_B(a_{ij})$ and b_{ij} by $-b_{ij}$ for all i, j , then we obtain $\partial b_{ij} = a_{ij} - \phi_B(a_{ij})$, which gives the braid DGA for B .

Next, as in [1], denote $\Delta = (\sigma_1 \cdots \sigma_{n-1})(\sigma_1 \cdots \sigma_{n-2}) \cdots (\sigma_1 \sigma_2) \sigma_1$. Then $\sigma_{n-k} = \Delta \sigma_k \Delta^{-1}$ for all k , and so \overline{B} and B^* are conjugate for any $B \in B_n$; hence by Theorem 2.7, the braid DGAs of \overline{B} and B^* are stable tame isomorphic.

It now suffices to show that B and B^* have equivalent braid DGAs. Consider the tame automorphism ξ of the algebra \mathcal{A}_n which sends a_{ij} to $a_{n+1-i, n+1-j}$ for each generator a_{ij} (and sends 1 to 1). By inspecting the definition of ϕ , we see that the induced action of ξ sends ϕ_{σ_k} to $\phi_{\sigma_{n-k}^{-1}}$, i.e., $\xi \phi_{\sigma_k} = \phi_{\sigma_{n-k}^{-1}} \xi$ for all k . Hence $\xi \phi_{B^*} = \phi_B \xi$ for all B .

Write the braid DGAs for B, B^* as $(\mathcal{A}, \partial), (\mathcal{A}, \tilde{\partial})$, respectively, and extend ξ to a tame automorphism of \mathcal{A} by sending b_{ij} to $b_{n+1-i, n+1-j}$ for each b_{ij} . Then

$$\partial \xi b_{ij} = \partial b_{n+1-i, n+1-j} = \xi(a_{ij}) - \phi_B(\xi(a_{ij})) = \xi(a_{ij} - \phi_{B^*}(a_{ij})) = \xi \tilde{\partial} b_{ij},$$

and so ξ intertwines the differentials ∂ and $\tilde{\partial}$. \square

Proposition 6.9. *The knot DGA classes for a knot, its mirror, and its inverse are the same.*

Proof. Let $B \in B_n$ be a braid whose closure is a knot K . We first show that the knot DGAs for B, B^* are stable tame isomorphic, which implies that K and its mirror have the same knot DGA class. Denote these knot DGAs by $(\mathcal{A}, \partial), (\mathcal{A}, \partial')$, respectively.

Let Ξ be the map on $n \times n$ matrices which permutes entries according to the definition $(\Xi(M))_{ij} = M_{n+1-i, n+1-j}$; note that Ξ commutes with matrix multiplication. Define the tame automorphism ξ on \mathcal{A} by its action on generators: $\xi(A) = \Xi(A)$, $\xi(B) = \Xi(B)$, $\xi(C) = \Xi(C)$, $\xi(D) = \Xi(D)$, $\xi(e_i) = e_{n+1-i}$.

Inspection of the definitions of ϕ and Φ^L shows that $\xi(\Phi_{\sigma_k}^L(A)) = \Xi(\Phi_{\sigma_{n-k}^L}^L(A))$ for all k . This, along with Proposition 4.4 and the identity $\xi\phi_{B^*} = \phi_B\xi$ from the proof of Proposition 6.8, allows us to prove readily by induction that $\xi(\Phi_{B^*}^L(A)) = \Xi(\Phi_B^L(A))$. Similarly, we can establish that $\xi(\Phi_{B^*}^R(A)) = \Xi(\Phi_B^R(A))$.

Hence

$$\partial\xi B = \Xi\partial B = (1 - \Xi(\Phi_B^L(A))) \cdot \Xi(A) = (1 - \xi(\Phi_{B^*}^L(A))) \cdot \xi(A) = \xi\partial' B$$

and similarly $\partial\xi C = \xi\partial' C$, while

$$\partial\xi D = \Xi\partial D = \xi(B) \cdot (1 - \xi(\Phi_{B^*}^R(A))) - (1 - \xi(\Phi_{B^*}^L(A))) \cdot \xi(C) = \xi\partial' D$$

and

$$\partial\xi e_i = \partial e_{n+1-i} = (\Xi(B + \Phi_B^L(A) \cdot C))_{ii} = (\xi(B) + \xi(\Phi_{B^*}^L(A)) \cdot \xi(C))_{ii} = \xi\partial' e_i.$$

It follows that ξ intertwines ∂ and ∂' , and so the knot DGAs of B and B^* are equivalent.

Now we show that the knot DGAs of B and B^{-1} are equivalent; this implies that K and its inverse have the same knot DGA class, since $B^{-1}, (B^{-1})^*$ have equivalent knot DGAs by the above argument, and $(B^{-1})^*, \overline{B^{-1}}$ are conjugate and hence have equivalent knot DGAs by Theorem 2.10. Write the knot DGA of B^{-1} as $(\tilde{\mathcal{A}}, \tilde{\partial})$, where the generators of $\tilde{\mathcal{A}}$ are marked by tildes. Identify the generators of $\tilde{\mathcal{A}}$ with those of \mathcal{A} by $\tilde{A} = \phi_B(A)$, $\tilde{B} = -B \cdot \Phi_B^R(A)$, $\tilde{C} = -\Phi_B^L(A) \cdot C$, $\tilde{D} = D$, and $\tilde{e}_i = d_{ii} - e_i$. Using Corollary 4.5 and Proposition 4.7, we calculate that

$$\partial\tilde{B} = (1 - \Phi_{B^{-1}}^L(\tilde{A})) \cdot \tilde{A} = (1 - (\Phi_B^L(A))^{-1}) \cdot \Phi_B^L(A) \cdot A \cdot \Phi_B^R(A) = \partial\tilde{B}$$

and similarly $\partial\tilde{C} = \partial\tilde{C}$, while

$$\partial\tilde{D} = \tilde{B} \cdot (1 - \Phi_{B^{-1}}^R(\tilde{A})) - (1 - \Phi_{B^{-1}}^L(\tilde{A})) \cdot \tilde{C} = B \cdot (1 - \Phi_B^R(A)) - (1 - \Phi_B^L(A)) \cdot C = \partial\tilde{D}$$

and

$$\partial\tilde{e}_i = (\tilde{B} + \Phi_{B^{-1}}^L(\tilde{A}) \cdot \tilde{C})_{ii} = (-B \cdot \Phi_B^R(A) - C)_{ii} = \partial(d_{ii} - e_i) = \partial\tilde{e}_i.$$

Hence our identification of generators, which yields a tame automorphism between \mathcal{A} and $\tilde{\mathcal{A}}$, gives $\partial = \tilde{\partial}$, and so the knot DGAs of B and B^{-1} are equivalent. \square

It can similarly be shown that the abelian braid and knot DGAs do not distinguish between mirrors or inverses. We may then ask how our invariants relate to classical invariants which also fail to distinguish between mirrors or inverses, such as the Alexander polynomial and the absolute value of the signature. In Section 7.1, we establish a connection between the knot DGA and the Alexander polynomial. First, in the next section, we deduce an invariant from the DGAs which shows that our invariants can distinguish between knots with the same Alexander polynomial and signature.

6.3. Augmentation number. We now introduce a family of easily computable invariants derived from the braid and knot DGAs. These invariants, which we call augmentation numbers, gives a crude but effective way to distinguish between braid conjugacy classes or knots. The advantage of augmentation numbers is that, unlike HC_0 or the full DGA, they are positive integers and hence require no ad hoc methods to interpret (cf. the computations of HC_0 in Section 8).

Definition 6.10. If R is a finitely generated ring and $d \geq 2$ an integer, define the \mathbb{Z}_d *augmentation number* of R to be the number of ring homomorphisms $R \rightarrow \mathbb{Z}_d$. For B any braid and K any knot, write $\text{Aug}(B, d)$, $\text{Aug}^{\text{ab}}(B, d)$, $\text{Aug}(K, d)$, $\text{Aug}^{\text{ab}}(K, d)$ for the \mathbb{Z}_d augmentation numbers of $HC_0(B)$, $HC_0^{\text{ab}}(B)$, $HC_0(K)$, $HC_0^{\text{ab}}(K)$, respectively.

We use the term ‘‘augmentation’’ in keeping with [8], in which the case $d = 2$ is discussed in the context of DGAs and Poincaré polynomials. One could imagine replacing \mathbb{Z}_d by any finite ring, although the usefulness of the more general construction is unclear.

Calculating augmentation numbers is straightforward. For instance, to compute $\text{Aug}(B, d)$ for $B \in B_n$, write $HC_0(B) = \mathbb{Z}\langle\{a_{ij}\}\rangle/\mathcal{I}$ and count the number of points in $\mathbb{Z}_d^{n(n-1)}$ (the space parametrizing ring homomorphisms $\mathbb{Z}\langle\{a_{ij}\}\rangle \rightarrow \mathbb{Z}_d$) which lie in the vanishing set of \mathcal{I} . There are $d^{n(n-1)}$ possible points and $n(n-1)$ conditions to check, corresponding to ∂b_{ij} for $1 \leq i \neq j \leq n$. Similarly, for $\text{Aug}^{\text{ab}}(B, d)$, there are $d^{n(n-1)/2}$ possible points and $n(n-1)/2$ conditions; for $\text{Aug}(K, d)$ with K the closure of B , $d^{n(n-1)}$ points and $2n^2$ conditions; and for $\text{Aug}^{\text{ab}}(K, d)$, $d^{n(n-1)/2}$ points and n^2 conditions. (Note thus that Aug^{ab} typically takes considerably less computing time than Aug to calculate.) In all cases, the \mathbb{Z}_d augmentation number is at least 1, as the following result shows.

Proposition 6.11. *For any braid B or knot K , the ring homomorphism from \mathcal{A}_n to \mathbb{Z} defined by sending each generator a_{ij} to -2 descends to homomorphisms from $HC_0(B)$, $HC_0^{\text{ab}}(B)$, $HC_0(K)$, $HC_0^{\text{ab}}(K)$ to \mathbb{Z} .*

Proof. It suffices to prove the assertion for $HC_0^{\text{ab}}(K)$, where K is the closure of the braid B ; see the diagram at the end of Section 6.1. Let M denote the $n \times n$ matrix all of whose entries are -2 ; then we need to show that $M = \Phi_B^L(M) \cdot M$. It is clear from the definition of ϕ and Lemma 4.6 that $\phi_{\sigma_k}(M) = M$ and $\Phi_{\sigma_k}^L(M) \cdot M = M$. By Proposition 4.4, it follows that

$$\Phi_{\sigma_k B}^L(M) \cdot M = \Phi_B^L(\phi_{\sigma_k}(M)) \cdot \Phi_{\sigma_k}^L(M) \cdot M = \Phi_B^L(M) \cdot M$$

for any B , and hence $\Phi_B^L(M) \cdot M = M$ for all B . \square

As an example of the computation of augmentation numbers, consider the trefoil 3_1 , for which we calculated in Section 4.1 that $HC_0 \cong \mathbb{Z}[x]/((x+2)(x-1))$; it is also easy to see that $HC_0^{\text{ab}}(3_1) \cong \mathbb{Z}[x]/((x+2)(x-1))$. Then $\text{Aug}(3_1, d) = \text{Aug}^{\text{ab}}(3_1, d) = 2$ for $d \neq 3$ and $\text{Aug}(3_1, 3) = \text{Aug}^{\text{ab}}(3_1, 3) = 1$. Note that these augmentation numbers could be calculated without a nice form for HC_0 , by directly using the definition of ∂ .

Any knot K with $HC_0(K) \cong \mathbb{Z}[x]/(p(x))$ for some polynomial $p(x)$ satisfies $\text{Aug}^{\text{ab}}(K, d) \leq \text{Aug}(K, d) \leq d$ for all d . Hence augmentation numbers can be used to show that some knots have HC_0 which is not of the form $\mathbb{Z}[x]/(p(x))$; see the examples in Sections 8.5 and 8.6.

7. KNOT CONTACT HOMOLOGY AND THE DETERMINANT

For knots K , contact homology is closely related to the classical invariant $|\Delta_K(-1)|$, the determinant of K . Throughout this section, we will denote by $\Sigma_2 = \Sigma_2(K)$ the double branched cover of S^3 over K , as in [22]; it is well known that $|H_1(\Sigma_2(K))| = |\Delta_K(-1)|$. In Section 7.1, we establish a surjection from $HC_0(K)$ to a quotient of the polynomial ring $\mathbb{Z}[x]$ specified by the largest invariant factor of $H_1(\Sigma_2(K))$, which typically is equal to the determinant of K . In Section 7.2, we discuss a natural linearization of knot contact homology, and see that $H_1(\Sigma_2(K))$, and hence the determinant, can be deduced from this linearization.

7.1. $HC_0(K)$ and $\Sigma_2(K)$. Before stating the main result of this section, we need to introduce a sequence of polynomials which plays a key role in the remainder of the paper. Define $\{p_m \in \mathbb{Z}[x]\}$ inductively by $p_0(x) = 2 - x$, $p_1(x) = x - 2$, $p_{m+1}(x) = xp_m(x) - p_{m-1}(x)$. Perhaps a more illuminating way to view p_m , $m \geq 1$, is via its roots:

$$p_m(x) = \prod_{k=0}^{m-1} \left(x - 2 \cos \frac{2k\pi}{2m-1} \right).$$

Theorem 7.1. *For K a knot, let $n(K)$ denote the largest invariant factor of the abelian group $H_1(\Sigma_2(K))$. Then there is a surjective ring homomorphism*

$$HC_0(K) \twoheadrightarrow \mathbb{Z}[x]/(p_{(n(K)+1)/2}(x)),$$

and this map factors through $HC_0^{\text{ab}}(K)$. In particular, the ranks of $HC_0(K)$ and $HC_0^{\text{ab}}(K)$ are at least $(n(K) + 1)/2$.

Note that $n(K)$ is odd since the determinant of a knot is always odd, and that $n(K)$ divides $|\Delta_K(-1)|$ and is divisible by any prime dividing $|\Delta_K(-1)|$. A table of invariant factors of $H_1(\Sigma_2(K))$ for small knots can be found in [2].

To prove Theorem 7.1, we will need a number of preliminary results. Define another sequence $\{q_m \in \mathbb{Z}[x]\}$ by $q_0(x) = -2$, $q_1(x) = -x$, $q_{m+1}(x) = xq_m(x) - q_{m-1}(x)$. Note that the recursion can be used to define q_m for all $m \in \mathbb{Z}$, and that $q_{-m} = q_m$ for all m because of symmetry.

Lemma 7.2. *For all $m_1, m_2 \in \mathbb{Z}$, we have*

$$q_{m_1}(x)q_{m_2}(x) + q_{m_1+m_2}(x) + q_{m_1-m_2}(x) = 0.$$

Proof. The identity holds by the defining relations for q_m for $m_2 = 0$ and $m_2 = 1$. It is then easy to establish by induction that it holds for all $m_2 \geq 0$, and hence for all m_2 since $q_{-m_2} = q_{m_2}$. \square

Lemma 7.3. *For all $\ell, m \in \mathbb{Z}$ with $m \geq 0$, we have $p_m(x) \mid q_{\ell+2m-1}(x) - q_\ell(x)$.*

Proof. An easy induction shows that $(p_m(x))^2 = (x+2)(q_{2m-1}(x)+2)$ and $p_m(x)p_{m+1}(x) = -(x+2)(q_{2m}(x)-x)$ for all $m \geq 0$; since $x+2$ divides $p_m(x)p_{m+1}(x)$, this establishes the lemma for $\ell = 0, 1$. The lemma now follows by induction on ℓ . \square

One can more elegantly prove Lemmas 7.2 and 7.3 by noting that the polynomials p_m, q_m are characterized by the identities $p_m(x+x^{-1}) = x^m + x^{-m} - x^{m-1} - x^{1-m}$, $q_m(x+x^{-1}) = -x^m - x^{-m}$; the lemmas then follow easily.

We next need to recall the Burau representation on B_n ; see [1] as a reference. For $1 \leq k \leq n-1$, define Bur_{σ_k} to be the linear map on the free $\mathbb{Z}[t, t^{-1}]$ module on n generators whose matrix is the identity except for the 2×2 submatrix formed by the $k, k+1$ rows and columns, which is $\begin{pmatrix} 1-t & t \\ 1 & 0 \end{pmatrix}$. This extends to a representation of B_n , classically known as the (nonreduced) *Burau representation*. The Burau representation is reducible to a trivial 1-dimensional representation spanned by the vector $(1, \dots, 1)$, and the $(n-1)$ -dimensional *reduced Burau representation*. More concretely, the map $\mathbb{Z}[t, t^{-1}]^n \rightarrow \mathbb{Z}[t, t^{-1}]^{n-1}$ given by $(v_1, \dots, v_n) \mapsto (v_2 - v_1, \dots, v_n - v_1)$ induces the reduced Burau representation on $\mathbb{Z}[t, t^{-1}]^{n-1}$.

For the remainder of this section, we will set $t = -1$; then the nonreduced and reduced Burau representations of B_n act on \mathbb{Z}^n and \mathbb{Z}^{n-1} , respectively. Let Bur_B and Bur_B^0 denote the matrices for B in the two respective representations. It is well known that Bur_B^0 is a presentation matrix for the group $H_1(\Sigma_2(K))$.

We need a bit more notation. If B is a braid, let \hat{B} denote the “reverse” braid obtained by reversing the word which gives B ; that is, if

$B = \sigma_{i_1}^{j_1} \sigma_{i_2}^{j_2} \cdots \sigma_{i_m}^{j_m}$, then $\hat{B} = \sigma_{i_m}^{j_m} \cdots \sigma_{i_2}^{j_2} \sigma_{i_1}^{j_1}$. (In our notation from Section 6.2, $\hat{B} = \overline{B}^{-1}$.) Also, for $v, w \in \mathbb{Z}^n$, let $M(v, w)$ denote the $n \times n$ matrix defined by $(M(v, w))_{ij} = q_{v_i - w_j}$.

Lemma 7.4. *Let $B \in B_n$ and $v \in \mathbb{Z}^n$. If we set $A = M(v, v)$, then $\phi_B(A) = M(\text{Bur}_{\hat{B}} v, \text{Bur}_{\hat{B}} v)$.*

Proof. Since $A = M(v, v)$, we have $a_{ij} = q_{v_i - v_j}$ for all i, j . The key is to verify the relation for $B = \sigma_k$. If $i \neq k, k+1$, then

$$\begin{aligned} \phi_{\sigma_k}(a_{ki}) &= -a_{k+1,i} - a_{k+1,k} a_{ki} = -q_{v_{k+1} - v_i} - q_{v_{k+1} - v_k} q_{v_k - v_i} = q_{2v_k - v_{k+1} - v_i} \\ &= (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{ki} \end{aligned}$$

by Lemma 7.2, and similarly $\phi_{\sigma_k}(a_{ik}) = (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{ik}$;

$$\phi_{\sigma_k}(a_{k+1,i}) = a_{ki} = q_{v_k - v_i} = (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{k+1,i}$$

and similarly $\phi_{\sigma_k}(a_{i,k+1}) = (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{i,k+1}$;

$$\phi_{\sigma_k}(a_{k,k+1}) = a_{k+1,k} = q_{v_{k+1} - v_k} = q_{v_k - v_{k+1}} = (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{k,k+1}$$

and similarly $\phi_{\sigma_k}(a_{k+1,k}) = (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{k+1,k}$; and finally, if $i, j \neq k, k+1$, then $\phi_{\sigma_k}(a_{ij}) = q_{v_i - v_j} = (M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v))_{ij}$. This shows that $\phi_{\sigma_k}(A) = M(\text{Bur}_{\sigma_k} v, \text{Bur}_{\sigma_k} v)$.

By replacing v by $\text{Bur}_{\sigma_k^{-1}} v$, we can deduce the desired identity for $B = \sigma_k^{-1}$ from the identity for $B = \sigma_k$. It thus suffices to show that if the identity holds for $B = B_1$ and $B = B_2$, then it holds for $B = B_1 B_2$. Now

$$\begin{aligned} \phi_{B_1 B_2}(M(v, v)) &= \phi_{B_1}(\phi_{B_2}(M(v, v))) \\ &= \phi_{B_2}(M(\text{Bur}_{\hat{B}_1} v, \text{Bur}_{\hat{B}_1} v)) \\ &= M(\text{Bur}_{\hat{B}_2} \text{Bur}_{\hat{B}_1} v, \text{Bur}_{\hat{B}_2} \text{Bur}_{\hat{B}_1} v) \\ &= M(\text{Bur}_{\widehat{B_1 B_2}} v, \text{Bur}_{\widehat{B_1 B_2}} v), \end{aligned}$$

and the lemma follows. \square

Lemma 7.5. *Let $B \in B_n$ and $v \in \mathbb{Z}^n$. If $A = M(v, v)$, then $\Phi_B^L(A) \cdot A = M(\text{Bur}_{\hat{B}} v, v)$ and $A \cdot \Phi_B^R(A) = M(v, \text{Bur}_{\hat{B}} v)$.*

Proof. We will prove the identity involving Φ^L ; the proof for Φ^R is completely analogous. First note that the identity is obvious when B is trivial. Now suppose that the identity holds for B ; we show that it holds for $B\sigma_k$. Indeed, by Proposition 4.4, we have

$$\Phi_{B\sigma_k}^L(A) \cdot A = \Phi_{\sigma_k}^L(\phi_B(A)) \cdot \Phi_B^L(A) \cdot A = \Phi_{\sigma_k}^L(\phi_B(A)) \cdot M(\text{Bur}_{\hat{B}} v, v).$$

By Lemma 4.6, $(\Phi_{B\sigma_k}^L(A) \cdot A)_{ij} = (\text{Bur}_{\hat{B}}v)_i - v_j$ except when $i = k$ or $i = k + 1$; in these cases, Lemmas 4.6, 7.2, and 7.4 yield

$$\begin{aligned} (\Phi_{B\sigma_k}^L(A) \cdot A)_{kj} &= -\phi_B(a_{k+1,k})(M(\text{Bur}_{\hat{B}}v, v))_{kj} - (M(\text{Bur}_{\hat{B}}v, v))_{k+1,j} \\ &= -(q(\text{Bur}_{\hat{B}}v)_{k+1} - (\text{Bur}_{\hat{B}}v)_k)(q(\text{Bur}_{\hat{B}}v)_{k-v_j}) - q(\text{Bur}_{\hat{B}}v)_{k+1-v_j} \\ &= q(2(\text{Bur}_{\hat{B}}v)_k - (\text{Bur}_{\hat{B}}v)_{k+1-v_j}), \end{aligned}$$

$$(\Phi_{B\sigma_k}^L(A) \cdot A)_{k+1,j} = (M(\text{Bur}_{\hat{B}}v, v))_{kj} = q(\text{Bur}_{\hat{B}}v)_{k-v_j}.$$

In all cases, we have $(\Phi_{B\sigma_k}^L(A) \cdot A)_{ij} = q(\text{Bur}_{\sigma_k \hat{B}}v)_{i-v_j}$, and so the identity holds for $B\sigma_k$. A similar computation shows that if the identity holds for B , then it holds for $B\sigma_k^{-1}$. The identity for all B follows by induction. \square

Proof of Theorem 7.1. Since $\text{Bur}_{\hat{B}}^0$ is a presentation matrix for $H_1(\Sigma_2(K)) = H_1(\Sigma_2(\bar{K}))$, and $n(K)$ is by definition an invariant factor of $H_1(\Sigma_2(K))$, there exists a vector $v = (v_2, \dots, v_n) \in \mathbb{Z}^{n-1}$ with $\gcd(v_2, \dots, v_n, n(K)) = 1$, such that the vector $(\text{Bur}_{\hat{B}}^0 - 1)v$ is divisible by $n(K)$. This means that $v' = (0, v_2, \dots, v_n) \in \mathbb{Z}^n$ satisfies the following property: if we write $w = (\text{Bur}_{\hat{B}} - 1)v'$, then $w_i \equiv w_j \pmod{n(K)}$ for all $1 \leq i, j \leq n$. Since the Burau representation sends $(1, \dots, 1)$ to itself, we may define $v'' = (-w_1, v_2 - w_1, \dots, v_n - w_1)$ and conclude that the vector $(\text{Bur}_{\hat{B}} - 1)v''$ is divisible by $n(K)$.

Now consider the ring homomorphism $\rho: \mathcal{A}_n \rightarrow \mathbb{Z}[x]$ defined by sending the matrix A to $M(v'', v'')$. By Lemma 7.5, ρ sends $(1 - \Phi_B^L(A)) \cdot A$ to $M(v'', v'') - M(\text{Bur}_{\hat{B}}v'', v'')$. Since $\text{Bur}_{\hat{B}}v'' \equiv v'' \pmod{n(K)}$, Lemma 7.3 shows that for all i, j ,

$$p_{(n(K)+1)/2} \mid q_{v''_i - v''_j} - q_{(\text{Bur}_{\hat{B}}v'')_{i-v''_j}} = (M(v'', v'') - M(\text{Bur}_{\hat{B}}v'', v''))_{ij}.$$

Hence $p_{(n(K)+1)/2}$ divides all entries of $\rho((1 - \Phi_B^L(A)) \cdot A)$. Similarly, $p_{(n(K)+1)/2}$ divides all entries of $\rho(A \cdot (1 - \Phi_B^R(A)))$. It follows that ρ descends to a map $\tilde{\rho}: HC_0(K) \rightarrow \mathbb{Z}[x]/(p_{(n(K)+1)/2}(x))$, which clearly factors through $HC_0^{\text{ab}}(K)$ since $q_m = q_{-m}$.

It remains to show that $\tilde{\rho}$ is surjective, or equivalently, that x is in the image of $\tilde{\rho}$. Define the set $S = \{j \in \mathbb{Z} \mid q_j(x) \in \text{im } \rho\}$. We first claim that $S = g\mathbb{Z}$ for some $g \in \mathbb{Z}$.

By Lemma 7.2, if $m_1 \in S$, then $m_1 + m_2 \in S$ if and only if $m_1 - m_2 \in S$. Hence if $a_1, a_2 \in S$, then the arithmetic progression $\{a_1 + \ell(a_2 - a_1)\}_{\ell \in \mathbb{Z}}$ is contained in S . Now suppose that $(m_1, m_2) = (a_1, a_2)$ minimizes $|m_1 - m_2|$, where (m_1, m_2) ranges over all pairs of distinct elements in S . Then $S = \{a_1 + \ell(a_2 - a_1)\}_{\ell \in \mathbb{Z}}$, since the existence of an element of S not in the arithmetic progression would violate minimality. Furthermore, $0 \in S$ since $q_0(x) = -2 \in \text{im } \rho$. Hence S is an ideal in \mathbb{Z} , which proves the claim.

Now for $2 \leq i \leq n$, we have $\rho(a_{i1}) = q_{v''_i - v''_1} = q_{v_i}$, and so $v_i \in S$. By the claim, we conclude that all multiples of $\gcd(v_2, \dots, v_n)$ are in S . Since $\gcd(v_2, \dots, v_n)$ and $n(K)$ are relatively prime by construction, there

exists $j \in S$ with $j \equiv 1 \pmod{n(K)}$. By Lemma 7.3, $p_{(n(K)+1)/2}$ divides $q_j(x) - q_1(x)$; but $q_j(x) \in \text{im } \rho$ and $q_1(x) = -x$, and hence $x \in \text{im } \tilde{\rho}$, as desired. \square

The above proof is a bit tedious and hard to digest. In [19], we will use the cord ring to see the surjection from Theorem 7.1 rather more directly, and justify the appearance of the polynomials p_m .

7.2. Linearized knot contact homology. Although it is generally difficult to find a nice form for the full knot contact homology $HC_*(K)$, there is a canonical linearization of $HC_*(K)$ which is easy to compute, and which encodes the group $H_1(\Sigma_2(K))$ and hence the determinant of K . We will discuss this linearization in this section.

Let (\mathcal{A}, ∂) be the knot DGA for a knot K . It follows easily from Proposition 6.11 that the algebra map $\epsilon: \mathcal{A} \rightarrow \mathbb{Z}$, sending each generator a_{ij} to -2 , and all other generators to 0, gives an augmentation for (\mathcal{A}, ∂) ; that is, $\epsilon \circ \partial = 0$. As in [3], we can construct linearized homology groups via this augmentation.

Let $\varphi_\epsilon: \mathcal{A} \rightarrow \mathcal{A}$ be the algebra isomorphism which sends a_{ij} to $a_{ij} - 2$ for all i, j , and acts as the identity on the other generators $b_{ij}, c_{ij}, d_{ij}, e_i$. Let \mathcal{M} be the subalgebra of \mathcal{A} generated by all words of length at least 1. Then $\partial_\epsilon := \varphi_\epsilon \circ \partial \circ \varphi_\epsilon^{-1}$ maps \mathcal{M} into itself. For any $k \geq 2$, ∂_ϵ descends to a differential $\partial_\epsilon^{\text{lin}}$ on the finite-rank graded \mathbb{Z} -module $\mathcal{M}/\mathcal{M}^k$, and the homology of the resulting complex gives a linearized version of $HC_*(K)$.

We now restrict our attention to the simplest nontrivial case, $k = 2$, although higher values of k also yield knot invariants. Note that $\mathcal{M}/\mathcal{M}^2$ is a free \mathbb{Z} -module generated in degree 0 by a_{ij} , in degree 1 by b_{ij} and c_{ij} , and in degree 2 by d_{ij} and e_i .

Definition 7.6. The *linearized contact homology* of K , written $HC_*^{\text{lin}}(K)$, is the graded homology of ∂_ϵ on $\mathcal{M}/\mathcal{M}^2$.

Using the conjugation symmetry of Section 6.1, we can also define an abelian version of linearized contact homology. In this case, (\mathcal{A}, ∂) is the abelian knot DGA of K with corresponding maximal subalgebra \mathcal{M} , and ∂_ϵ descends to a differential on $\mathcal{M}/\mathcal{M}^2$, whose homology we denote by $HC_*^{\text{ab,lin}}(K)$.

Proposition 7.7. *The groups $HC_*^{\text{lin}}(K)$ and $HC_*^{\text{ab,lin}}(K)$ are invariants of the knot K .*

Proof. All of the stable tame isomorphisms used in the proof of Theorem 2.10 commute with ϵ , because $\phi_B \circ \epsilon = \epsilon \circ \phi_B$ for any braid B . The proposition follows. \square

We now establish some results about $HC_*^{\text{lin}}(K)$. Note that this homology exists only in dimensions $* = 0, 1, 2$.

Proposition 7.8. *For K a knot, $HC_2^{\text{lin}}(K) \cong \mathbb{Z}$ and $HC_1^{\text{lin}}(K) \cong (H_1(\Sigma_2) \oplus \mathbb{Z}) \otimes (H_1(\Sigma_2) \oplus \mathbb{Z}) \oplus \mathbb{Z}^m$ for some $m \geq 0$.*

Proposition 7.8 will follow almost immediately from a lemma which we now formulate. Let d_1, d_2, \dots, d_k be the invariant factors of $H_1(\Sigma_2)$, with $d_1 | d_2 | \dots | d_k$. Write $Y = \text{Bur}_{\hat{B}}$, and express $1 - Y$ in Smith normal form; more precisely, there are matrices $X_1, X_2 \in GL(n, \mathbb{Z})$ with $X_1 \cdot (1 - Y) X_2 = \Delta$, where Δ is a diagonal matrix. Since $1 - Y$ is a presentation matrix for $H_1(\Sigma_2) \oplus \mathbb{Z}$, we may assume that the diagonal entries of Δ are precisely d_1, \dots, d_n , where $d_{k+1} = \dots = d_{n-1} = 1$ and $d_n = 0$.

Now write the linearized chain complex of the alternate knot DGA of the braid B as $C_2 \xrightarrow{M_2} C_1 \xrightarrow{M_1} C_0$, where C_0, C_1, C_2 are generated by $\{a_{ij}\}, \{b_{ij}, c_{ij}, d_{ij}\}, \{e_{ij}, f_{ij}\}$, respectively. The map M_2 is given by the action of $\partial_\epsilon^{\text{lin}}$ on C_2 : $\partial_\epsilon^{\text{lin}} E = B - D - C \cdot (\text{Bur}_{\hat{B}})^T$, $\partial_\epsilon^{\text{lin}} F = B - C - \text{Bur}_{\hat{B}} \cdot D$.

Lemma 7.9. *After changing bases for C_1 and C_2 , we can write M_2 as a diagonal $(3n^2 - n) \times (2n^2)$ matrix, with n^2 diagonal entries given by $\gcd(d_i, d_j) = d_{\min(i,j)}$, $1 \leq i, j \leq n$, and the remaining n^2 diagonal entries given by 1.*

Proof. We wish to calculate Smith normal form for M_2 . Replace D by $D + B - C \cdot Y^T$ and then F by $F + Y \cdot E$ (this simply reparametrizes d_{ij} and f_{ij}); then $\partial_\epsilon^{\text{lin}} E = -D$ and $\partial_\epsilon^{\text{lin}} F = B - C - Y \cdot (B - C \cdot Y^T)$. The action of $\partial_\epsilon^{\text{lin}}$ on E gives n^2 “1” diagonal entries in M_2 . To complete the Smith normal form for M_2 , we need to examine the matrix $Z = B - C - Y \cdot (B - C \cdot Y^T)$.

Introduce dummy variables b_{11}, \dots, b_{nn} (this adds n rows of zeros to M_2), and replace b_{ij} by $b_{ij} - b_{jj}$ for all $i \neq j$. If we write $\tilde{B} = (b_{ij})$, which is identical to B away from the diagonal, then Z becomes $\tilde{B} - C - Y \cdot (\tilde{B} - C \cdot Y^T)$. Next replace Z by $X_1 \cdot Z \cdot X_1^T$; this has the effect of changing basis in C_2 . Finally, change basis in C_1 by successively replacing \tilde{B} by $X_2 \cdot \tilde{B} \cdot (X_1^T)^{-1} + C$, and then C by $Y^{-1} \cdot X_1^{-1} \cdot C \cdot X_2^T$. Then Z becomes

$$\Delta \cdot \tilde{B} - C \cdot \Delta = (d_i b_{ij} - d_j c_{ij}).$$

The lemma follows immediately. \square

Proof of Proposition 7.8. We have $HC_2^{\text{lin}}(K) = \ker M_2$ and $HC_1^{\text{lin}}(K) = (\ker M_1) / (\text{im } M_2)$. Write M_2 as in Lemma 7.9. Since $d_1, \dots, d_{n-1} > 0$, M_2 has only one 0 diagonal entry, and so $\ker M_2 \cong \mathbb{Z}$.

For $HC_1^{\text{lin}}(K)$, assume that we have chosen bases for C_1, C_2 so that M_2 is diagonal with diagonal entries $\delta_1, \dots, \delta_{2n^2}$, with $\delta_{2n^2} = 0$ and $\delta_i \neq 0$ for $i < 2n^2$. Let the corresponding basis for C_1 be v_1, \dots, v_{3n^2-n} . Then $v_1, \dots, v_{2n^2-1} \in \ker M_1$, and indeed we may assume that $\{v_1, \dots, v_{2n^2-1+m}\}$ forms a basis for $\ker M_1$ for some $m \geq 0$. It follows that $(\ker M_1) / (\text{im } M_2) \cong \mathbb{Z}^m \oplus \oplus_i \mathbb{Z}_{\delta_i}$. From Lemma 7.9, we conclude that $(\ker M_1) / (\text{im } M_2) \cong (H_1(\Sigma_2) \oplus \mathbb{Z}) \otimes (H_1(\Sigma_2) \oplus \mathbb{Z}) \oplus \mathbb{Z}^m$, as desired. \square

We conjecture that m is always 0 in Proposition 7.8. Even if this is not the case, Proposition 7.8 still implies that we can deduce $H_1(\Sigma_2)$ from $HC_1^{\text{lin}}(K)$; this is an easy consequence of the fundamental theorem of finitely generated abelian groups.

Corollary 7.10. *The group $H_1(\Sigma_2)$, and hence the determinant $|\Delta_K(-1)|$, is determined by the equivalence class of the knot DGA of K .*

We now turn our attention to $HC_0^{\text{lin}}(K)$, and derive a result which will be used in the sequel [19].

Proposition 7.11. *There are surjections*

$$HC_0^{\text{lin}}(K) \twoheadrightarrow HC_0^{\text{lin,ab}}(K) \twoheadrightarrow \text{Sym}^2(H_1(\Sigma_2)).$$

The proof of Proposition 7.11 is essentially a linearized version of the (more involved) proof of Theorem 7.1, and we provide just an outline here. The matrix $\text{Bur}_{\hat{B}}$ acts on \mathbb{Z}^n , which has generators e_1, \dots, e_n , and it is an easy exercise to check that $H_1(\Sigma_2)$ is generated by the differences $\{e_i - e_j\}$, modulo the relations $\text{Bur}_{\hat{B}}e_i - e_i = 0$ for all i . (Note that $\text{Bur}_{\hat{B}}e_i - e_i$ is a linear combination of differences $e_{j_1} - e_{j_2}$.)

Lemma 7.12. *In the ring $\mathbb{Z}[e_1, \dots, e_n]$, if $A = (-2 + (e_i - e_j)^2)$, then*

$$\Phi_B^L(A) \cdot A = (-2 + (\text{Bur}_{\hat{B}}e_i - e_j)^2 + O(e^4))$$

and

$$\Phi_B^R(A) \cdot A = (-2 + (e_i - \text{Bur}_{\hat{B}}e_j)^2 + O(e^4)).$$

Proof. Induction on the braid word B , as in the proof of Lemma 7.5. \square

Proof of Proposition 7.11. The surjection from $HC_0^{\text{lin}}(K)$ to $\text{Sym}^2(H_1(\Sigma_2))$, which factors through $HC_0^{\text{lin,ab}}(K)$, sends a_{ij} to $(e_i - e_j)^2$. This is well-defined by Lemma 7.12 and the fact that $\text{Bur}_{\hat{B}}e_i - e_i = 0$ in $H_1(\Sigma_2)$ for all i ; it is a surjection by polarization, since 2 is invertible in $\text{Sym}^2(H_1(\Sigma_2))$. \square

A more natural way to see the second surjection in Proposition 7.11, using cords, is given in Section 5.2 of [19]. By all computational indications, it seems likely that this surjection is always an isomorphism, and hence that $HC_0^{\text{lin,ab}}(K)$ is determined by $H_1(\Sigma_2)$ for all knots K . However, $HC_0^{\text{lin}}(K)$ is not determined by $H_1(\Sigma_2)$. For instance, the knots 8_{18} and 9_{37} both satisfy $H_1(\Sigma_2) \cong \mathbb{Z}_3 \oplus \mathbb{Z}_{15}$, but $HC_0^{\text{lin}}(8_{18}) \cong \mathbb{Z}_3^3 \oplus \mathbb{Z}_{15}$, while $HC_0^{\text{lin}}(9_{37}) \cong \mathbb{Z}_3^2 \oplus \mathbb{Z}_{15}$. It is possible that $HC_0^{\text{lin}}(K)$ is always isomorphic to either $\text{Sym}^2(H_1(\Sigma_2))$ or $H_1(\Sigma_2) \otimes H_1(\Sigma_2)$; this has been verified for all prime knots with eleven or fewer crossings.

8. COMPUTATIONS

The combinatorial nature of our invariants allows them to be calculated readily by computer. *Mathematica* source to compute the invariants is available at the author's web site. The full program requires the installation of

the noncommutative algebra package NCAAlgebra/NCGB [11], though there is a commutative version which does not use NCAAlgebra. Using the program, we can calculate the braid homomorphism ϕ , braid and knot DGAs, HC_0 (via Gröbner bases), augmentation numbers, and linearized knot contact homology. Many of the calculations in this section were performed with the help of the program.

8.1. Braid conjugacy classes. It can be difficult to tell if two braids have equivalent braid DGAs. Currently, the most effective computational tool in this regard is the augmentation number invariant described in Section 6.3, which can be calculated by computer and is reasonably effective in distinguishing between braid conjugacy classes.

In Table 1, we list all conjugacy classes in B_3 of nonnegative writhe which can be represented by a connected braid of word length at most 6, along with their writhe and selected augmentation numbers. (Recall that the writhe w is the homomorphism $B_n \rightarrow \mathbb{Z}$ for which $w(\sigma_k) = 1$ for all k .) The augmentation numbers, combined with the writhe, distinguish all of the given conjugacy classes, with the exception of the mirrors $\sigma_1^2\sigma_2^{-2}\sigma_1\sigma_2^{-1}$ and $\sigma_1^{-2}\sigma_2^2\sigma_1^{-1}\sigma_2$, which by Proposition 6.8 cannot be distinguished using our invariants.

Without the use of the writhe, our invariants become less effective. For instance, it can be shown that the braid DGAs for $\sigma_1\sigma_2$ and $\sigma_1^3\sigma_2$ are both

B	w	$\text{Aug}(B, 2)$	$\text{Aug}^{\text{ab}}(B, 2)$	$\text{Aug}^{\text{ab}}(B, 5)$	$\text{Aug}^{\text{ab}}(B, 7)$
$\sigma_1^5\sigma_2$	6	5	1	4	6
$\sigma_1^3\sigma_2^3$	6	9	5	12	18
$\sigma_1^3\sigma_2$	4	4	2	5	7
$\sigma_1^5\sigma_2^{-1}$	4	4	2	5	6
$\sigma_1^3\sigma_2\sigma_1^{-1}\sigma_2$	4	4	2	4	7
$\sigma_1\sigma_2$	2	4	2	5	7
$\sigma_1^3\sigma_2^{-1}$	2	4	2	4	7
$\sigma_1^3\sigma_2^{-1}\sigma_1\sigma_2^{-1}$	2	2	2	9	4
$\sigma_1\sigma_2^{-1}$	0	5	1	4	6
$\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}$	0	9	5	12	18
$\sigma_1^3\sigma_2^{-3}$	0	7	5	11	13
$\sigma_1^2\sigma_2^{-2}\sigma_1\sigma_2^{-1}$	0	1	1	4	2
$\sigma_1^{-2}\sigma_2^2\sigma_1^{-1}\sigma_2$	0	1	1	4	2

TABLE 1. The conjugacy classes in B_3 with nonnegative writhe which can be represented by a connected braid of word length at most 6, identified by such a representative B . The writhe w is given, along with various augmentation numbers.

stable tame isomorphic to the DGA on two generators of degree 0 and two generators of degree 1, with $\partial = 0$.

8.2. The unknot and unlink, and other links. The unknot is the closure of the trivial braid in B_1 , whose knot DGA is generated by b_{11}, c_{11} in degree 1 and d_{11}, e_1 in degree 2, with differential given by $\partial b_{11} = \partial c_{11} = \partial d_{11} = 0$, $\partial e_1 = b_{11} + c_{11}$. (The 1×1 matrices Φ^L and Φ^R for the trivial braid in B_1 are simply (1).) This is stable tame isomorphic to the DGA with generators b_{11} in degree 1 and d_{11} in degree 2, and with $\partial = 0$. In particular, we see that $HC_0(\text{unknot}) \cong \mathbb{Z}$.

More generally, let L_n denote the n -component unlink, which is the closure of the trivial braid B_n^0 in B_n . The matrices $\Phi_{B_n^0}^L$ and $\Phi_{B_n^0}^R$ are both the identity, and so the knot DGA of B_n^0 has differential $\partial a_{ij} = \partial b_{ij} = \partial c_{ij} = \partial d_{ij} = 0$, $\partial e_i = b_{ii} + c_{ii}$, and $HC_0(L_n) \cong \mathbb{Z}\langle\{a_{ij}\}\rangle$.

By contrast, the Hopf link, which is the closure of $\sigma_1^2 \in B_2$, has knot DGA satisfying $\partial B = \begin{pmatrix} -4+a_{12}a_{21} & 4a_{12}-a_{12}a_{21}a_{12} \\ 0 & -4+a_{21}a_{12} \end{pmatrix}$ and $\partial C = \begin{pmatrix} -4+a_{12}a_{21} & 0 \\ 4a_{21}-a_{21}a_{12}a_{21} & -4+a_{21}a_{12} \end{pmatrix}$. It follows that $HC_0(\text{Hopf}) \cong \mathbb{Z}\langle a_{12}, a_{21} \rangle / \langle -4 + a_{12}a_{21}, -4 + a_{21}a_{12} \rangle$, which is not isomorphic to $HC_0(L_2)$.

Finally, consider the split link $3_1 \sqcup 0_1$ given by the closure of $\sigma_1^3 \in B_3$, which is the unlinked union of an unknot and a trefoil. We can compute that

$$HC_0(3_1 \sqcup 0_1) \cong \mathbb{Z}\langle a_{12}, a_{13}, a_{31}, a_{23}, a_{32} \rangle / \langle -2 + a_{12} + a_{12}^2, \\ a_{13} - a_{23} - a_{12}a_{13} + a_{12}a_{23}, a_{31} - a_{32} - a_{31}a_{12} + a_{32}a_{12} \rangle,$$

which is not isomorphic to $HC_0(L_2)$. Hence our invariants, in contrast to the Alexander polynomial, can distinguish between split links.

8.3. Torus knots $T(2, n)$. In this section, we generalize our computation in Section 4.1 of HC_0 for the trefoil, by computing HC_0 for all torus knots of the form $T(2, n)$.

First we introduce two sequences related to the polynomials $q_m(x)$ discussed in Section 7.1. In \mathcal{A}_2 , define $\{q_m^{(1)}\}, \{q_m^{(2)}\}$ by $q_0^{(1)} = q_0^{(2)} = -2$, $q_1^{(1)} = a_{12}$, $q_1^{(2)} = a_{21}$, and $q_{m+1}^{(1)} = -a_{12}q_m^{(2)} - q_{m-1}^{(1)}$, $q_{m+1}^{(2)} = -a_{21}q_m^{(1)} - q_{m-1}^{(2)}$; by definition, we have $q_m^{(1)}|_{a_{12}=a_{21}=-x} = q_m^{(2)}|_{a_{12}=a_{21}=-x} = q_m(x)$.

Lemma 8.1. *For $k \geq 1$, we have*

$$\Phi_{\sigma_1^{2k-1}}^L(A) \cdot A = \begin{pmatrix} q_{2k-1}^{(2)} & q_{2k}^{(2)} \\ q_{2k-2}^{(1)} & q_{2k-1}^{(1)} \end{pmatrix} \quad \text{and} \quad A \cdot \Phi_{\sigma_1^{2k-1}}^R(A) = \begin{pmatrix} q_{2k-1}^{(1)} & q_{2k-2}^{(1)} \\ q_{2k}^{(2)} & q_{2k-1}^{(2)} \end{pmatrix}.$$

Proof. We prove the first identity; the second follows similarly or by conjugation symmetry. For $k = 1$, the identity can be verified directly. For $k \geq 2$, by Proposition 4.4, we have

$$\Phi_{\sigma_1^{2k-1}}^L(A) \cdot A = \Phi_{\sigma_1^2}^L(\phi_{\sigma_1^{2k-3}}(A)) \cdot \Phi_{\sigma_1^{2k-3}}^L(A) \cdot A.$$

Now $\phi_{\sigma_1^2}(A) = A$, and so $\phi_{\sigma_1^{2m-1}}(A) = \phi_{\sigma_1}(A) = \begin{pmatrix} -2 & a_{21} \\ a_{12} & -2 \end{pmatrix}$; then we compute that $\Phi_{\sigma_1^2}^L(\phi_{\sigma_1^{2m-1}}(A)) = \begin{pmatrix} -1+a_{21}a_{12} & a_{21} \\ -a_{12} & -1 \end{pmatrix}$. The desired identity can now be checked by induction on m . \square

We can now compute HC_0 for the $(2, 2k - 1)$ torus knot.

Proposition 8.2. *For $k \geq 1$, $HC_0(T(2, 2k - 1)) \cong \mathbb{Z}[x]/(p_k(x))$, where p_k is the polynomial defined in Section 7.1.*

Proof. By Lemma 8.1, we have $((1 - \Phi_{\sigma_1^{2k-1}}^L(A)) \cdot A)_{12} = a_{12} - q_{2k}^{(2)}$ and $(A \cdot (1 - \Phi_{\sigma_1^{2k-1}}^R(A)))_{21} = a_{21} - q_{2k}^{(2)}$. Hence in $HC_0(T(2, 2k - 1))$, we have $a_{12} = a_{21}$, and we can set $x := -a_{12} = -a_{21}$. The relations in $HC_0(T(2, 2k - 1))$ then become the entries of the matrix $\begin{pmatrix} -2 - q_{2k-1}(x) & -x - q_{2k}(x) \\ -x - q_{2k-2}(x) & -2 - q_{2k-1}(x) \end{pmatrix}$.

Now define $r_m(x) = q_m(x) - q_{2k-1-m}(x)$ for all m , so that $r_m(x) = xr_{m+1}(x) - r_{m+2}(x)$. We have

$$\begin{aligned} HC_0(T(2, 2k - 1)) &\cong \mathbb{Z}[x]/(-x - q_{2k-2}(x), -x - q_{2k}(x), -2 - q_{2k-1}(x)) \\ &= \mathbb{Z}[x]/(r_0(x), r_1(x)), \end{aligned}$$

where the second equality holds because $-x - q_{2k}(x) = r_{-1}(x) = xr_0(x) - r_1(x)$. Now

$\gcd(r_0(x), r_1(x)) = \gcd(r_1(x), r_2(x)) = \cdots = \gcd(r_{k-1}(x), r_k(x)) = r_{k-1}(x)$;
since $r_{k-1}(x) = p_k(x)$ by induction on k , the proposition follows. \square

Proposition 8.2 shows that the surjection in Theorem 7.1 is actually an isomorphism for $(2, n)$ torus knots. This is established in more generality, for all two-bridge knots, in [19]. However, we will see in the following sections that it is not true for all knots.

Before moving on, we make an observation regarding the trefoil in the context of Legendrian knot theory. We have seen that HC_0 for the trefoil is $\mathbb{Z}[x]/(x^2 + x - 2)$. This has two maps to \mathbb{Z}_2 and thus the knot DGA for the trefoil has two augmentations over \mathbb{Z}_2 . If we use the form for the trefoil knot DGA given at the end of Section 2.2, then one augmentation, ϵ_0 , sends a_{12} and a_{21} to 0, while the other, ϵ_1 , sends them to 1. The linearized differential for ϵ_0 is simply given by the linear terms in the differential mod 2, and one can calculate that this has Poincaré polynomial $\lambda^2 + \lambda$ (i.e., homology of dimension 1 in degrees 2 and 1). On the other hand, one can check that the augmentation ϵ_1 yields Poincaré polynomial $4\lambda^2 + 4\lambda$. This is an example where different augmentations of a DGA measuring relative contact homology give different Poincaré polynomials over \mathbb{Z}_2 ; this phenomenon has also now been observed for some Legendrian knots in \mathbb{R}^3 [17].

8.4. Torus knots $T(3, n)$. The invariant HC_0 is not as simple for general torus knots as it is for $T(2, n)$. For instance, if we write the torus knot

$T(3, 4)$ as the closure of the braid $(\sigma_1\sigma_2)^4 \in B_3$, we can calculate

$$\begin{aligned} HC_0(T(3, 4)) \cong \mathbb{Z}\langle a_1, a_2 \rangle / \langle & 2a_1 - 2a_2 - a_1^2 + a_1a_2 + a_1^3 - a_1^2a_2, \\ & -a_1a_2 + a_2a_1 + a_1^2a_2 - a_2a_1^2, -3a_1 + 3a_2 + a_1a_2a_1 - a_2a_1a_2, \\ & -a_1 + a_2 + a_1^2 - a_2^2, 2 + a_2 - a_1a_2 - a_2a_1 - a_1^3 \rangle. \end{aligned}$$

Furthermore, if we use lexicographic order with $a_1 < a_2$, then the above presentation gives a Gröbner basis for the ideal. Thus as an additive abelian group, $HC_0(T(3, 4)) \cong \mathbb{Z}^7$, generated by $1, a_1, a_2, a_1^2, a_1a_2, a_2a_1, a_1a_2a_1$. On the other hand, the abelianization of $HC_0(T(3, 4))$ is easily shown to be isomorphic to $\mathbb{Z}[x]/(x^3 - 2x^2 - x + 2)$, and so $HC_0(T(3, 4))$ is noncommutative. Note that $HC_0(T(3, 4))$ does map surjectively to $\mathbb{Z}[x]/(x^2 - x - 2)$, as per Theorem 7.1.

In general, it seems that $HC_0(T(m, n))$ is too complicated to be expressed in closed form. However, for $T(3, n)$, there is a relatively simple expression for HC_0^{ab} , if not for HC_0 (cf. $HC_0(T(3, 4))$ above).

It is easy to see that $HC_0^{\text{ab}}(T(3, n))$ can be written in the form $\mathbb{Z}[x]/(p(x))$ for some polynomial p . The knot $T(3, n)$ is the closure of $(\sigma_1\sigma_2)^n \in B_3$, and $\phi_{(\sigma_1\sigma_2)^n}$ cyclically permutes a_{12}, a_{23}, a_{31} when 3 does not divide n . Hence the generators a_{12}, a_{23}, a_{13} of $HC_0^{\text{ab}}(T(3, n))$ are equal by Corollary 4.9. (The same argument shows that $HC_0(T(3, n))$ is generated by two elements.)

Explicitly calculating $HC_0^{\text{ab}}(T(3, n))$ is more involved; a straightforward but tedious induction, along the lines of the proof of Proposition 8.2, yields the following result.

Proposition 8.3. *If $\gcd(3, n) = 1$, then*

$$HC_0^{\text{ab}}(T(3, n)) \cong \begin{cases} \mathbb{Z}[x]/(r_k(x)), & n = 2k - 1 \\ \mathbb{Z}[x]/(s_k(x)), & n = 2k - 2, \end{cases}$$

where $r_0(x) = 2 - x$, $r_1(x) = x - 2$, $r_{m+1}(x) = (x - 1)r_m(x) - r_{m-1}(x)$, and $s_1(x) = 0$, $s_2(x) = (x - 2)(x + 1)$, $s_{m+1}(x) = (x - 1)s_m(x) - s_{m-1}(x)$.

Note that $r_k(x) = \frac{x-2}{x-3} p_k(x - 1)$.

8.5. The connected sums $3_1\#3_1$ and $3_1\#\overline{3_1}$. When we view the connected sum $3_1\#3_1$ as the closure of $\sigma_1^3\sigma_2^3 \in B_3$, we can calculate that $\text{Aug}^{\text{ab}}(3_1\#3_1, 2) = 7$, and hence $HC_0(3_1\#3_1)$ is not of the form $\mathbb{Z}[x]/(p(x))$. To garner more information, we use the standard Gröbner basis algorithm to find that

$$\begin{aligned} HC_0(3_1\#3_1) \cong \mathbb{Z}\langle a_1, a_2, a_3, a_4 \rangle / \langle & (a_1 + 2)(a_1 - 1), (a_2 + 2)(a_2 - 1), \\ & (a_1 - 1)a_3, a_4(a_1 - 1), (a_2 - 1)a_4, a_3(a_2 - 1) \rangle. \end{aligned}$$

The given presentation for the above ideal is a noncommutative Gröbner basis for the ideal with respect to any lexicographic order, and it follows that $HC_0(3_1\#3_1)$ is noncommutative (for instance, $a_1a_3 \neq a_3a_1$) and not

finitely generated as a \mathbb{Z} -module. It is also true that HC_0^{ab} has infinite rank in this case; in fact,

$$HC_0^{\text{ab}}(3_1\#3_1) \cong \mathbb{Z}[a_1, a_2, a_3]/((a_1 + 2)(a_1 - 1), (a_2 + 2)(a_2 - 1), (a_1 - 1)(a_3 - a_2), (a_2 - 1)(a_3 - a_1)).$$

As another application of this example, it turns out that precisely the same computation holds for $3_1\#\overline{3_1}$; hence it is not true that HC_0 or HC_0^{ab} determines the isotopy class of a knot, even up to mirrors.

8.6. Knot contact homology and other invariants. From the computations of the previous sections, the HC_0 invariant for knots depends on more than the determinant of the knot. It is also not determined by the Alexander polynomial; for instance, the knots 6_1 and 9_{46} have the same Alexander polynomial, but $\text{Aug}^{\text{ab}}(6_1, 2) = 2$ while $\text{Aug}^{\text{ab}}(9_{46}, 2) = 5$.

More generally, knot contact homology seems to behave independently of many known knot invariants. In particular, HC_0 can distinguish knots which share many of the same “classical” invariants.

The knots 10_{25} and $\overline{10_{56}}$ (the mirror of 10_{56} , with the conventions of [22]) are both alternating, with the same HOMFLY polynomial and signature; hence they also share the same Alexander polynomial, Jones polynomial, Khovanov invariant [13] (by a result of [15]), and Ozsváth–Szabó invariant [21]. On the other hand, we have $\text{Aug}^{\text{ab}}(10_{25}, 7) = 1$, while $\text{Aug}^{\text{ab}}(\overline{10_{56}}, 7) = 2$.

Similarly, the knots 11_{255}^a and 11_{257}^a , alternating 11-crossing knots from Conway’s enumeration [4], have the same (two-variable) Kauffman polynomial, but $\text{Aug}^{\text{ab}}(11_{255}^a, 7) = 1$ while $\text{Aug}^{\text{ab}}(11_{257}^a, 7) = 2$. We conclude the following result.

Proposition 8.4. *The invariant HC_0 for knots is not determined by any of the following: Alexander polynomial, Jones polynomial, HOMFLY polynomial, Kauffman polynomial, signature, Khovanov invariant, and Ozsváth–Szabó invariant.*

9. AN ALTERNATE KNOT INVARIANT

As explained in Section 3, in the symplectic world, the knot DGA counts holomorphic disks with certain boundary conditions determined by the knot. This count is well defined modulo 2, and so we obtain a DGA over \mathbb{Z}_2 . To lift this DGA to \mathbb{Z} coefficients, however, we need a set of coherent orientations for the relevant moduli spaces. The knot DGA we have been using presumably corresponds to a choice of coherent orientations, but there are others.

By experimenting with various choices of signs, one can derive several inequivalent choices for DGAs over \mathbb{Z} which are knot invariants and agree with the original knot DGA modulo 2, but only two seem to give “nice” results for simple knots (e.g., such that HC_0 for small knots is of the form

$\mathbb{Z}[x]/(p(x))$). One is the knot DGA already described; the other we call the *alternate knot DGA*. This alternate theory lacks some of the nice properties of the knot DGA—for instance, no obvious analogue of Theorem 7.1 exists for the alternate DGA—but it seems to give a sharper knot invariant than the usual DGA.

9.1. The alternate DGA. We follow the definition of the knot DGA, making adjustments where necessary.

Let $\tilde{\phi}$ denote the homomorphism from B_n to $\text{Aut}(\mathcal{A}_n)$ defined on generators by

$$\tilde{\phi}_{\sigma_k} : \begin{cases} a_{ki} & \mapsto -a_{k+1,i} + a_{k+1,k}a_{ki} & i \neq k, k+1 \\ a_{ik} & \mapsto -a_{i,k+1} - a_{ik}a_{k,k+1} & i \neq k, k+1 \\ a_{k+1,i} & \mapsto a_{ki} & i \neq k, k+1 \\ a_{i,k+1} & \mapsto a_{ik} & i \neq k, k+1 \\ a_{k,k+1} & \mapsto -a_{k+1,k} \\ a_{k+1,k} & \mapsto -a_{k,k+1} \\ a_{ij} & \mapsto a_{ij} & i, j \neq k, k+1. \end{cases}$$

Note that $\tilde{\phi}$ is identical to ϕ modulo 2, and that ϕ and $\tilde{\phi}$ are conjugate through the automorphism of \mathcal{A}_n which sends a_{ij} to a_{ij} if $i < j$ and $-a_{ij}$ if $i > j$.

Write $A = (a_{ij})$, $B = (b_{ij})$, $C = (c_{ij})$, $D = (d_{ij})$ as usual, with the crucial difference that we set $a_{ii} = 0$, rather than $a_{ii} = -2$, for all i . Denote by $\tilde{\phi}^{\text{ext}}$ the composition map $B_n \hookrightarrow B_{n+1} \xrightarrow{\tilde{\phi}} \text{Aut}(\mathcal{A}_n)$, and define matrices $\tilde{\Phi}_B^L(A)$, $\tilde{\Phi}_B^R(A)$ by

$$\tilde{\phi}_B^{\text{ext}}(a_{i*}) = \sum_{j=1}^n (\tilde{\Phi}_B^L(A))_{ij} a_{j*} \quad \text{and} \quad \tilde{\phi}_B^{\text{ext}}(a_{*j}) = \sum_{i=1}^n a_{*i} (\tilde{\Phi}_B^R(A))_{ij}.$$

Then the *alternate knot DGA* of $B \in B_n$ is generated, as before, by $\{a_{ij} \mid 1 \leq i \neq j \leq n\}$ in degree 0, $\{b_{ij}, c_{ij} \mid 1 \leq i, j \leq n\}$ in degree 1, and $\{d_{ij} \mid 1 \leq i, j \leq n\}$ and $\{e_i \mid 1 \leq i \leq n\}$ in degree 2, with

$$\begin{aligned} \partial A &= 0 \\ \partial B &= (1 + \tilde{\Phi}_B^L(A)) \cdot A \\ \partial C &= A \cdot (1 + \tilde{\Phi}_B^R(A)) \\ \partial D &= B \cdot (1 + \tilde{\Phi}_B^R(A)) - (1 + \tilde{\Phi}_B^L(A)) \cdot C \\ \partial e_i &= (B - \tilde{\Phi}_B^L(A) \cdot C)_{ii}. \end{aligned}$$

Note the difference in signs from Definition 2.9. Also, let the *alternate contact homology* of a knot K , written $\widetilde{HC}_*(K)$, be the homology of the alternate knot DGA of any braid closing to K . As mentioned before, the alternate knot DGA and contact homology are identical modulo 2 to the usual knot DGA and contact homology.

Theorem 9.1. *The stable tame isomorphism class of the alternate knot DGA, as well as \widetilde{HC}_* , are knot invariants.*

The proof of Theorem 9.1 is entirely analogous to the proofs from Sections 4 and 5, and is omitted here.

Many of the results from Section 6 still hold in this context. We can define conjugation on the alternate knot DGA by $\overline{a_{ij}} = -a_{ji}$, $\overline{b_{ij}} = -c_{ji}$, $\overline{c_{ij}} = -b_{ji}$, $\overline{d_{ij}} = d_{ji}$, and $\overline{e_i} = d_{ii} - e_i$; then the differential ∂ in the alternate knot DGA commutes with conjugation. Hence we can form the *alternate abelian knot DGA* similarly to Definition 6.5, but with $a_{ji} = -a_{ij}$ and $c_{ij} = -b_{ji}$. None of the alternate invariants distinguish between knot mirrors or inverses.

9.2. Computations of $\widetilde{HC}_0(K)$ for small knots. One interesting question is whether an analogue of Theorem 7.1 holds for \widetilde{HC}_0 . In all computations performed by the author, there is a surjection from $\widetilde{HC}_0(K)$ to some ring of the form $\mathbb{Z}[x]/(p(x))$, where p is monic of degree $\frac{n(K)+1}{2}$ and $p(x) \equiv p_{(n(K)+1)/2}(x) \pmod{2}$. Why this should be true in general is unclear.

Nevertheless, for small knots K , we have $\widetilde{HC}_0(K) \cong \mathbb{Z}[x]/(p(x))$ for some polynomial p with $\deg(p) = (|\Delta_K(-1)| + 1)/2$. In contrast to the case of HC_0 , however, the polynomial p depends on more than just $|\Delta_K(-1)|$. For example, the knots 4_1 and 5_1 both satisfy $|\Delta_K(-1)| = 5$ and $HC_0(K) \cong \mathbb{Z}[x]/(x^3 - x^2 - 3x + 2)$, but $\widetilde{HC}_0(4_1) \cong \mathbb{Z}[x]/(x^3 - x^2 + x)$, while $\widetilde{HC}_0(5_1) \cong \mathbb{Z}[x]/(x^3 - x^2 - x)$. See Table 2 for more examples.

Proposition 9.2. *If K is a prime knot with seven or fewer crossings, then we can write $\widetilde{HC}_0(K) \cong \mathbb{Z}[x]/(p(x))$ with $\deg(p) = (|\Delta_K(-1)| + 1)/2$. Furthermore, p can be expressed as $p = \tilde{p}_{(|\Delta_K(-1)|+1)/2}$ for some sequence $\{\tilde{p}_m\}$ with $\tilde{p}_0(x) = \tilde{p}_1(x) = x$ and $\tilde{p}_{m+1}(x) = x\tilde{p}_m(x) \pm \tilde{p}_{m-1}(x)$ for each m ; in particular, $p \equiv p_{(|\Delta_K(-1)|+1)/2} \pmod{2}$, with $\{p_m\}$ as in Section 7.1.*

Proposition 9.2 can be verified by direct computation using the author's program in *Mathematica*; see Table 2.

As with HC_0 , it is easy to compute \widetilde{HC}_0 explicitly for any torus knot $T(2, n)$.

Proposition 9.3. *For $k \geq 1$, $HC_0(T(2, 2k - 1)) \cong \mathbb{Z}[x]/(\tilde{p}_k(x))$, where*

$$\tilde{p}_k(x) = \sum_{j=0}^{k-1} (-1)^{\lfloor j/2 \rfloor} \binom{k-1-\lfloor j/2 \rfloor}{\lfloor j/2 \rfloor} x^{k-j}.$$

Note that the sequence $\{\tilde{p}_k\}$ in Proposition 9.3 satisfies $\tilde{p}_0(x) = \tilde{p}_1(x) = x$ and $\tilde{p}_{m+1}(x) = x\tilde{p}_m(x) - \tilde{p}_{m-1}(x)$, and that $\tilde{p}_m(x) = xp_m(x)/(x+2)$ for all m .

In general, when $\widetilde{HC}_0(K)$ is of the form $\mathbb{Z}[x]/(p(x))$ with p monic, the polynomial $p(x)$ is not unique; however, we can deduce two useful invariants of the isomorphism class of $\widetilde{HC}_0(K)$ from $p(x)$. The first is the degree of p ,

K	$p(x)$ with $\widetilde{HC}_0(K) \cong \mathbb{Z}[x]/(p(x))$	$D(p)$
0_1	x	1
3_1	$x^2 - x$	1
4_1	$x^3 - x^2 + x$	-3
5_1	$x^3 - x^2 - x$	5
5_2	$x^4 - x^3 + x$	-23
6_1	$x^5 - x^4 + x^3 + x$	257
6_2	$x^6 - x^5 - 2x^4 + x^3 + x^2 + x$	1777
6_3	$x^7 - x^6 - x^5 + 2x^4 - x^2 + x$	-10571
7_1	$x^4 - x^3 - 2x^2 + x$	49
7_2	$x^6 - x^5 + x^3 + x^2 - x$	4409
7_3	$x^7 - x^6 - 3x^5 + 2x^4 + 2x^3 + x^2 - x$	78301
7_4	$x^8 - x^7 + 4x^6 - 3x^5 + 4x^4 - 2x^3 + x$	-1166607
7_5	$x^9 - x^8 - x^7 + 2x^6 + x^5 - 2x^4 + 2x^2 - x$	-4690927
7_6	$x^{10} - x^9 + 2x^8 - x^7 + 3x^6 - x^5 + 2x^4 + x^2 + x$	90320393
7_7	$x^{11} - x^{10} + 3x^9 - 2x^8 + 4x^7 - 3x^6 + 3x^5 - 2x^4 + x^3 - x^2 + x$	-932501627

TABLE 2. The invariant $\widetilde{HC}_0(K)$ for prime knots with seven or fewer crossings, and the discriminant $D(p)$. Knot notation is as in [22].

which is simply the vector space dimension of $\widetilde{HC}_0(K; \mathbb{Q})$. The second is the discriminant $D(p)$, the well-known integer invariant which can be defined as $\prod_{i < j} (r_i - r_j)^2$, where $\{r_i\}$ is the set of roots of p .

Table 2 gives $\widetilde{HC}_0(K)$ for all prime knots K with seven or fewer crossings, while Table 3 contrasts the invariants HC_0 and \widetilde{HC}_0 . Note that the discriminant on \widetilde{HC}_0 allows us to distinguish between knots which have the same HC_0 ; in other words, lifting $\widetilde{HC}_0(K; \mathbb{Z}_2) = HC_0(K; \mathbb{Z}_2)$ to $\widetilde{HC}_0(K)$ does give new information. This would seem to provide an instance where considering orientations in contact homology yields results not given by the unoriented theory.

REFERENCES

- [1] J. Birman, *Braids, Links, and Mapping Class Groups*, Ann. of Math. Studies **82** (Princeton University Press, Princeton, 1974).
- [2] G. Burde and H. Zieschang, *Knots*, 2nd ed., de Gruyter Studies in Mathematics **5** (Walter de Gruyter & Co., Berlin, 2003).
- [3] Yu. Chekanov, Differential algebra of Legendrian links, *Invent. Math.* **150** (2002), no. 3, 441–483.
- [4] J. H. Conway, An enumeration of knots and links, and some of their algebraic properties, in *Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)*, pp. 329–358 (Pergamon, Oxford, 1970).

K	$p(x)$ with $HC_0(K) \cong \mathbb{Z}[x]/(p(x))$	$p(x)$ with $\widetilde{HC}_0(K) \cong \mathbb{Z}[x]/(p(x))$
0_1	$x - 2$	x
3_1	$x^2 - x - 2$	$x^2 - x$
4_1	$x^3 - x^2 - 3x + 2$	$x^3 - x^2 + x$
5_1	$x^3 - x^2 - 3x + 2$	$x^3 - x^2 - x$
5_2	$x^4 - x^3 - 4x^2 + 3x + 2$	$x^4 - x^3 + x$
6_1	$x^5 - x^4 - 5x^3 + 4x^2 + 5x - 2$	$x^5 - x^4 + x^3 + x$
6_2	$x^6 - x^5 - 6x^4 + 5x^3 + 9x^2 - 5x - 2$	$x^6 - x^5 - 2x^4 + x^3 + x^2 + x$
6_3	$x^7 - x^6 - 7x^5 + 6x^4 + 14x^3 - 9x^2 - 7x + 2$	$x^7 - x^6 - x^5 + 2x^4 - x^2 + x$
7_1	$x^4 - x^3 - 4x^2 + 3x + 2$	$x^4 - x^3 - 2x^2 + x$
7_2	$x^6 - x^5 - 6x^4 + 5x^3 + 9x^2 - 5x - 2$	$x^6 - x^5 + x^3 + x^2 - x$
7_3	$x^7 - x^6 - 7x^5 + 6x^4 + 14x^3 - 9x^2 - 7x + 2$	$x^7 - x^6 - 3x^5 + 2x^4 + 2x^3 + x^2 - x$

TABLE 3. Comparison between HC_0 and \widetilde{HC}_0 for some small knots.

- [5] T. Ekholm, J. Etnyre, and M. Sullivan, Legendrian submanifolds in R^{2n+1} and contact homology, preprint, [arXiv:math.SG/0210124](#).
- [6] Ya. Eliashberg, Invariants in contact topology, *Doc. Math.* **1998**, Extra Vol. II, 327–338 (electronic).
- [7] Ya. Eliashberg, A. Givental, and H. Hofer, Introduction to Symplectic Field Theory, *GAF A 2000 (Tel Aviv, 1999)*, *Geom. Funct. Anal.* **2000**, Special Volume, Part II, 560–673.
- [8] J. Epstein, D. Fuchs, and M. Meyer, Chekanov–Eliashberg invariants and transverse approximations of Legendrian knots, *Pacific J. Math.* **201** (2001), no. 1, 89–106.
- [9] J. Etnyre, L. Ng, and J. Sabloff, Invariants of Legendrian knots and coherent orientations, *J. Symplectic Geom.* **1** (2002), no. 2, 321–367.
- [10] K. Fukaya and Y.-G. Oh, Zero-loop open strings in the cotangent bundle and Morse homotopy, *Asian J. Math.* **1** (1997), no. 1, 96–180.
- [11] J. Helton, R. Miller, and M. Stankus, NCAAlgebra: a Mathematica package for doing non commuting algebra, <http://www.math.ucsd.edu/~ncalg>.
- [12] S. Humphries, An approach to automorphisms of free groups and braids via transvections, *Math. Z.* **209** (1992), no. 1, 131–152.
- [13] M. Khovanov, A categorification of the Jones polynomial, *Duke Math. J.* **101** (2000), no. 3, 359–426.
- [14] D. Kramer, The braid group B_4 is linear, *Invent. Math.* **142** (2000), no. 3, 451–486.
- [15] E. S. Lee, On Khovanov invariant for alternating links, preprint, [arXiv:math.GT/0210213](#).
- [16] W. Magnus, Rings of Fricke characters and automorphism groups of free groups, *Math. Z.* **170** (1980), no. 1, 91–103.
- [17] P. Melvin and S. Shrestha, The nonuniqueness of Chekanov polynomials of Legendrian knots, preprint, [arXiv:math.GT/0411206](#).
- [18] L. Ng, Computable Legendrian invariants, *Topology* **42** (2003), no. 1, 55–82.
- [19] L. Ng, Knot and braid invariants from contact homology II, preprint, [arXiv:math.GT/0303343](#).
- [20] H. Ooguri and C. Vafa, Knot invariants and topological strings, *Nuclear Phys. B* **577** (2000), no. 3, 419–438.

- [21] P. Ozsváth and Z. Szabó, Holomorphic disks and knot invariants, *Adv. Math.* **186** (2004), no. 1, 58–116.
- [22] D. Rolfsen, *Knots and Links* (Publish or Perish Press, Berkeley, 1976).
- [23] J. Sabloff, Invariants of Legendrian knots in circle bundles, *Commun. Contemp. Math.* **5** (2003), no. 4, 569–627.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305
E-mail address: `lng@math.stanford.edu`